

**A new Plethystic Symmetric Function Operator
and
The rational Compositional Shuffle Conjecture at $t=1/q$
by**

A.M. Garsia, E. Leven, N. Wallach and G. Xin

Abstract. Our main result here is that the specialization at $t = 1/q$ of the $Q_{km, kn}$ operators studied in [4] may be given a very simple plethystic form. This discovery yields elementary and direct derivations of several identities relating these operators at $t = 1/q$ to the Rational Compositional Shuffle conjecture of [3]. In particular we show that if m, n and k are positive integers and (m, n) is a coprime pair then

$$q^{\frac{(km-1)(kn-1)+k-1}{2}} Q_{km, kn}(-1)^{kn} \Big|_{t=1/q} = \frac{[k]_q}{[km]_q} e_{km} [X[km]_q] \quad \text{A.1}$$

where as customarily, for any integer $s \geq 0$ and indeterminate u we set $[s]_u = 1 + u + \cdots + u^{s-1}$. We also show that the symmetric polynomial on the right hand side is always Schur positive. Moreover, using the Rational Compositional Shuffle conjecture, we derive a precise formula expressing this polynomial in terms of Parking functions in the $km \times kn$ lattice rectangle.

Introduction

The specializations at $t = 1/q$ of all the Shuffle conjectures (including the classical cases) are still open to this date. What makes this specialization particularly fascinating is that both sides of the stated identities have combinatorial interpretations. Nevertheless proving these identities is quite challenging even in the simplest cases. For instance from the Rational Shuffle Conjecture we can easily derive the following identity, for any coprime pair (m, n) .

$$\sum_{D \in \mathcal{D}_{m,n}} q^{\text{coarea}(D) + \text{dinv}(D)} = \frac{1}{[m]_q} \begin{bmatrix} m+n-1 \\ n \end{bmatrix}_q \quad \text{I.1}$$

where the sum is over Dyck paths in the $m \times n$ lattice rectangle, $\text{coarea}(D)$ gives the number of lattice squares above the path and $\text{dinv}(D)$ is a Dyck path statistic that can also be given a relatively simple geometric construction. The identity obtained by setting $q = 1$ in I.1 is an immediate consequence of the Cyclic Lemma, which suggests that this classical result may have a natural q -analogue. The investigations that yielded the present results have been directed towards the necessity of giving a concrete setting to a variety of identities stated or implied in recent work by the Algebraic Geometers, most particularly in the papers Burban-Schiffmann [5] and Schiffmann-Vasserot [27]. Unfortunately most of this work appears in language that requires considerable algebraic Geometrical background. We have been privileged to have had the statements of some of these results translated in a language that we could understand by Eugene Gorsky and Andrey Negut. Many of the theorems we prove here have their origin in this algebraic geometrical literature. Our contribution is to provide proofs that are accessible to the algebraic combinatorial audience. We hope that in doing so, the new results we obtain may be conducive to progress in this most challenging area of Algebraic Combinatorics.

We will be dealing here with an algebra $\mathcal{A}$ of linear operators acting on the space Λ of symmetric function in an infinite alphabet $X = \{x_1, x_2, x_3, \dots\}$ with coefficients in the field $\mathbb{Q}(q, t)$ of rational functions in the two indeterminates q, t . Given a symmetric function $F[X] \in \Lambda$, it will be convenient to denote by “ $\underline{F}$ ” the operator “multiplication by $F[X]$ ”. As customary we will denote by “ $F^\perp$ ” the operator dual of $\underline{F}$ with respect to the classical Hall scalar product of symmetric functions.

For a coprime pair (m, n) the $Q_{m,n}$ operators have an elementary definition, which as far as we understand, is due to Burban-Schiffmann in [5]. By taking a lattice point (a, b) in the $m \times n$ rectangle that is closest and below the segment $(0, 0) \leftrightarrow (m, n)$ and setting $(c, d) = (m, n) - (a, b)$ we obtain a decomposition

$$(m, n) = (a, b) + (c, d) \quad \text{I.2}$$

which here and after will be referred to as “*Split*”(m, n). Since the co-primality of the pair (m, n) forces both pairs (a, b) and (b, c) to be coprime, we can recursively set

$$Q_{m,n} = \frac{1}{M} [Q_{c,d}, Q_{a,b}], \quad (M = (1-t)(1-q)) \quad \text{I.3}$$

with base cases

$$Q_{0,1} = -e_1 \quad \text{and} \quad Q_{1,0} = D_0 \quad \text{I.4}$$

where, e_1 is the customary elementary symmetric function and D_0 belongs to a family of operators $\{D_k\}_{k \in \mathbf{Z}}$, introduced in [12], and defined by setting for $F[X] \in \Lambda$

$$D_k F[X] = F[X + \frac{M}{z}] \sum_{r \geq 0} (-z)^r e_r[X] \Big|_{z^k} \quad \text{I.5}$$

In dealing with the present subject it is absolutely indispensable to make use of plethystic notation. Readers that are not familiar with this device are referred to [14] for an introduction to its use. The readers will also find in [3] and [4], elementary proofs of all the auxiliary results that we will need in this writing. Those papers were written precisely to render this subject accessible to the algebraic combinatorial audience in a completely self contained manner. In particular it is shown in [4] that to compute the action of an operator $Q_{m,n}$ we do not need to recurse to the base cases in I.4, but rather use as a short cut the identities

$$Q_{1,k} = D_k \quad \text{I.6}$$

It should be mentioned that the original identities justifying the use of this short cut were first given in [12].

The definition of the operator $Q_{u,v}$ for a non coprime pair (u, v) relies on a truly amazing property of the algebra generated by the operators D_k , certainly noticed in [5] and possibly in other algebraic geometrical literature. To state it it will be convenient to write a non-coprime pair in the form $(u, v) = (km, kn)$ with (m, n) coprime and $k > 1$. This given, we can recursively define the operator $Q_{km,kn}$ by choosing any of the lattice points (a, b) in the rectangle $km \times kn$ that are strictly below and closest to the segment $(0, 0) \leftrightarrow (km, kn)$ then set

$$Q_{km,kn} = \frac{1}{M} [Q_{km-a, kn-b}, Q_{a,b}]. \quad \text{I.7}$$

This definition is made possible because, the choice of (a, b) forces both (a, b) and $(km - a, kn - b)$ to be coprime. Moreover, all the operators resulting from such a choice of (a, b) can be shown to act identically on symmetric functions.

Another fundamental fact discovered by the Algebraic Geometers is that the Q operators indexed by collinear vectors do commute. More precisely for any coprime pair m, n and any two integers k, h we have

$$[Q_{km,kn}, Q_{hm,hn}] = 0. \quad \text{I.8}$$

Elementary, but by no means simple, proofs of all these properties are given in [4]. The complexity of these proofs is due to the recursive nature of the definition in I.3. Our discovery here is that with the specialization $t = 1/q$ we can obtain several explicit identities from which these fundamental properties are immediate.

More precisely let $D_{u,v}$ denote the operator whose action on the symmetric function $F[X] \in \Lambda$ is defined by setting

$$D_{u,v}F[X] = F[X + M[u]_q/z] \sum_{r \geq 0} (-z)^r e_r[[u]_t X] \Big|_{z^v}, \quad \text{I.9}$$

where we must set here $t = 1/q$.

Theorem I.1

If a, b, c, d, u, v are any integers related by the vector identity $(a, b) + (c, d) = (u, v)$ we have for non vanishing a, c, u

$$\frac{1}{M} [D_{c,d}, D_{a,b}] \Big|_{t=1/q} = q^{1+bc} \frac{[a]_q [c]_q}{[u]_q} \frac{1 - q^{da-bc}}{1 - q} D_{u,v} \Big|_{t=1/q} \quad \text{I.11}$$

This identity has the following immediate corollary

Theorem I.2

For any coprime pair (m, n) and $k \geq 1$ we have

$$q^{(km-1)(kn-1)/2+(k-1)/2} Q_{km, kn} \Big|_{t=1/q} = q^{(km-1)kn} \frac{[k]_q}{[km]_q} D_{km, kn}. \quad \text{I.12}$$

The two identities in I.11 and I.12 have variety of consequences. For instance we can immediately see from I.11 that the collinearity of (a, b) and (c, d) implies that $D_{a,b}$ and $D_{c,d}$ commute. We thus obtain a much simpler proof of this commutativity result for the $Q_{u,v}$ operators when $t = 1/q$. Another immediate consequence of I.11 is that the algebra generated by the D_k operators at $t = 1/q$ is spanned by the *convex* monomials in the $D_{u,v}$ operators. Here a monomial

$$D_{u_1, v_1} D_{u_2, v_2} D_{u_3, v_3} \cdots D_{u_l, v_l},$$

is called *convex* if and only if we have

$$\frac{v_1}{u_1} \geq \frac{v_2}{u_2} \geq \cdots \geq \frac{v_l}{u_l}.$$

To state an important consequence of I.12, we need some background. Let us recall that the classical Shuffle conjecture of Haglund et Al in [20] may be stated as the identity

$$Q_{n+1, n}(-1)^n = \sum_{PF \in \mathcal{PF}_n} t^{\text{area}(PF)} q^{\text{dinv}(PF)} s_{\text{pides}(PF)}[X] \quad \text{I.13}$$

where the sum is over Parking Functions in the $n \times n$ lattice square, $\text{area}(PF)$ and $\text{dinv}(PF)$ are Parking function statistics we will define later in a much more general context, and $s_{\text{pides}(PF)}[X]$ denotes the Schur function indexed by the composition which gives the descent set of a permutation naturally associated to a Parking Function. In a recent paper E. Gorsky and A. Negut formulated an infinite variety Shuffle conjectures, one for each coprime pair (m, n) . They may be stated in a form similar to I.13, namely

$$Q_{m, n}(-1)^n = \sum_{PF \in \mathcal{PF}_{m, n}} t^{\text{area}(PF)} q^{\text{dinv}(PF)} s_{\text{pides}(PF)}[X] \quad \text{I.14}$$

where the sum is over Parking functions in the $m \times n$ lattice rectangle, and the parking function statistics occurring in I.14 are highly non trivial modifications of the statistics involved in I.12. Now, Theorem I.2 has the following immediate corollary

Theorem I.3

For any coprime pair (m, n) and $k \geq 1$ we have

$$q^{(km-1)(kn-1)/2+(k-1)/2} Q_{km, kn}(-1)^{kn} \Big|_{t=1/q} = \frac{[k]_q}{[km]_q} e_{kn}[X[km]_q]. \quad \text{I.15}$$

In particular, by combining I.15 with the Gorsky-Negut conjectures at $t = 1/q$ we obtain the identity

$$\frac{1}{[m]_q} e_n[X[m]_q] = \sum_{PF \in \mathcal{PF}_{m,n}} q^{\text{coarea}(PF) + \text{dinu}(PF)} s_{\text{pides}(PF)}[X] \quad \text{I.16}$$

with $\text{coarea}(PF) = (m-1)(n-1)/2 - \text{area}(PF)$.

This given, we may ask if the right hand side of I.15, can also be given a Parking Function interpretation. It turns out that this indeed the case. More precisely we will show the following

Theorem I.4

Upon the validity of the extended Compositional Shuffle Conjecture in [3] it follows that

$$\frac{[k]_q}{[km]_q} e_{kn}[X[km]_q] = \sum_{PF \in \mathcal{PF}_{km, kn}} q^{\text{coarea}(PF) + \text{dinu}(PF)} [\text{ret}(PF)]_q s_{\text{pides}(PF)}[X] \quad \text{I.17}$$

where $\text{ret}(PF)$ is a statistic which indicates the height of the first return to the diagonal by the Dyck path of PF in the $km \times kn$ lattice rectangle.

The precise definitions of all the Parking Function statistics occurring in I.17 will be given in the sequel.

We must mention that it would follow from I.17 combined with the theory of LLT polynomials that the left hand side is a Schur positive symmetric polynomial. However, we will show that this particular result can be given a much more elementary proof.

It is important to notice that operators $D_{u,v}$ can be used for any integral values of u and v . Now it follows from I.12 for $m = 1$ and $n = 0$ that

$$Q_{k,0} \Big|_{t=1/q} = D_{k,0} \quad \text{I.18}$$

It was known to the Algebraic Geometers that the family of operators $\{Q_{k,0}\}_{k \geq 1}$ have as a complete set of eigenfunctions the modified Macdonald basis $\{\tilde{H}_\mu[X; q, t]\}_\mu$ introduced in [11]. More precisely, we have

$$Q_{k,0} \tilde{H}_\mu[X; q, t] = \left(1 - (1 - t^k)(1 - q^k) B_\mu(q^k, t^k)\right) \tilde{H}_\mu[X; q, t], \quad \text{I.19}$$

where for a partition $\mu = (\mu_1, \mu_2, \dots, \mu_l)$ we set $B_\mu(q, t) = \sum_{i=1}^l t^{i-1} \sum_{j=1}^{\mu_i} q^{j-1}$. Since it can be shown that the polynomial $\tilde{H}_\mu[X; q, t]$ specializes, at $t = 1/q$ to a scalar multiple of $s_\mu[\frac{X}{1-q}]$, it follows from I.19 and I.18 that we must also have

Theorem I.5

$$D_{k,0} s_\mu\left[\frac{X}{1-q}\right] = \left(1 - (1 - q^{-k})(1 - q^k) B_\mu(q^k, q^{-k})\right) s_\mu\left[\frac{X}{1-q}\right] \quad \text{I.20}$$

Proving this identity directly from the definition in I.9 leads to some highly non trivial combinatorial problems. However, with some effort, as we shall see, a less direct but still entirely elementary path to I.20 can actually be found. In fact this particular effort led to the discovery of the following formula for the action of $D_{u,v}$ on the basis $\{s_\mu[\frac{X}{1-q}]\}_\mu$.

Theorem I.6

For any $u, v > 0$ and any partition μ we have

$$D_{u,v} s_\mu\left[\frac{X}{1-q}\right] = (q^u - 1) \sum_{i=1}^{|\mu|+v} q^{up(\mu)_i + v - ui} s_{p(\mu) + ve_i}\left[\frac{X}{1-q}\right] \quad \text{I.21}$$

where $p(\mu)$ is the weak composition of length $|\mu| + v$ obtained by adjoining zeros to the parts of μ and e_i is the i^{th} coordinate vector of length $|\mu| + v$.

Remark I.1

We should mention that there is another interesting by-product of our introduction of the operator $D_{u,v}$. We learned from Eugene Gorsky (see also section 6.10 of [16]) that in [29] it is shown that for a suitable constant factor $c_{m,n}(q)$ we have, for (m, n) a coprime pair

$$Q_{m,n} \Big|_{t=1/q} = c_{m,n}(q) \nabla^{\frac{m}{n}} p_n \nabla^{-\frac{m}{n}} \Big|_{t=1/q} \quad \text{I.22}$$

with ∇ the operator introduced in [2]. Now it turns out that one can easily derive I.22 from I.21 for $(u, v) = (m, n)$, directly from the definition of ∇ given in [2].

This paper is divided into five sections. In the first section we give an elementary proof of Theorem I.1. This type of proof has been successfully used in various similar situations where we needed a straightforward proof of an identity that was discovered by another path. In this section we also give proofs of Theorems I.2 and I.3.

In section 2 we give the elementary argument that proves the polynomiality and Schur positivity of the symmetric function in I.15.

In section 3 we give our Parking Function setting for the symmetric function in I.15. We also give there a simplified version of the Parking Function statistics that occur in the formulation of the Rational Compositional Shuffle Conjecture that take account of the most recent developments in this subject.

In section 4 we prove Theorems I.5 and I.6 and explore some of their consequences. What is interesting is that the path to these proofs uses an argument that may be conducive to the discovery of a variety other identities of similar type.

In section 5 we give an outline of the theoretical steps that led to the discovery of the operators $D_{u,v}$ and Theorem I.1.

Acknowledgement

We cannot overemphasize the importance of the contributions to the very existence of this article by Eugene Gorsky and Andrey Negut. The situation we find ourselves in is that a variety of symmetric function constructs and combinatorial conjectures that have been discovered and extensively studied by the Algebraic Combinatorists for more than 2 decades has been recently enriched in a truly spectacular fashion by discoveries of the Algebraic Geometers, (see for instance [5], [16], [17], [18], [9], [24], [26], [27], [28], [29]). Unfortunately most of this information is written in a language that is accessible only to people with extensive Algebraic Geometrical background. Over a period of nearly two years Gorsky and Negut provided us with extensive information about these developments in a language that we could understand. The present article is now the fourth in a series of articles ([3], [4], [9]) where our principal aim has been to give these contributions of the Algebraic Geometers a self contained Algebraic Combinatorial setting, in the hope that this enrichment of our subject will be conducive to further fruitful Algebraic Combinatorial developments.

1. Commutator properties of our new operators

Our main goal in this section is an elementary proof of Theorem I.1 and its immediate Corollaries. In a later section we will try to give a glimpse of the machinery that led to the discovery of the operators $D_{u,v}$ and yielded the original proof of this identity. To give the reader an idea of the basic difference between these two approaches, we need to recall a device which was extensively used in all previous work in the theory of Macdonald polynomials. We will refer to it as the “ Ω ” notation. The point of departure of plethystic substitutions is the operation of evaluating the power symmetric function p_k at a formal power series $E = E(t_1, t_2, t_3, \dots)$ containing an unlimited number of indeterminates. We simply set

$$p_k[E] = E(t_1^k, t_2^k, t_3^k, \dots) \quad 1.1$$

Since every symmetric function can be expressed as a polynomial in the power functions, this definition allows to evaluate $F[E]$ for any given symmetric function F . This is what we refer to as the plethystic substitution of E in F . In this vein we set

$$\Omega[E] = \exp\left(\sum_{k \geq 1} \frac{p_k[E]}{k}\right). \quad 1.2$$

Clearly, this definition implies that for any two expressions A and B we have

$$\Omega[A + B] = \Omega[A] \times \Omega[B] \quad \text{and} \quad \Omega[A - B] = \Omega[A]/\Omega[B] \quad 1.3$$

In particular it also follows from 1.1 and 1.2, that if $X = x_1 + x_2 + x_3 + \dots$ then

$$\sum_{r \geq 0} (-z)^r e_r[X] = \Omega[-zX]. \quad 1.4$$

Using this device the definition of the operators D_k in I.5 can be rewritten in the form

$$D_k F[X] = F[X + \frac{M}{z}] \Omega[-zX] \Big|_{z^k} \quad 1.5$$

Successive applications of two operators D_a and D_b to a symmetric function $F[X]$, in this notation, leads to the identities

$$\begin{aligned} D_b D_a F[X] &= D_b F[X + \frac{M}{z_1}] \Omega[-z_1 X] \Big|_{z_1^a} \\ &= F[X + \frac{M}{z_1} + \frac{M}{z_2}] \Omega[-z_1(X + \frac{M}{z_2})] \Omega[-z_2 X] \Big|_{z_1^a z_2^b} \\ &= F[X + \frac{M}{z_1} + \frac{M}{z_2}] \Omega[-(z_1 + z_2)X] \Omega[-\frac{Mz_1}{z_2}] \Big|_{z_1^a z_2^b} \\ &= F[X + \frac{M}{z_1} + \frac{M}{z_2}] \Omega[-(z_1 + z_2)X] \frac{(1-z_1/z_2)(1-qtz_1/z_2)}{(1-tz_1/z_2)(1-qz_1/z_2)} \frac{1}{z_1^a z_2^b} \Big|_{z_1^0 z_2^0} \end{aligned} \quad 1.6$$

where the last equality results from multiple applications of the identities in 1.2. By contrast if we carry out this calculation, the way a computer would do it, we would end up with the following sequence of identities.

$$\begin{aligned} D_b D_a F[X] &= D_b \sum_{r_1 \geq 0} F^{(r_1)}[X] \frac{1}{z_1^{r_1}} \Omega[-z_1 X] \Big|_{z_1^a} \\ &= D_b \sum_{r_1 \geq 0} F^{(r_1)}[X] (-1)^{r_1+a} e_{r_1+a}[X] \\ &= \sum_{r_1 \geq 0} F^{(r_1)}[X + \frac{M}{z_2}] (-1)^{r_1+a} e_{r_1+a}[X + \frac{M}{z_2}] \Omega[-z_2 X] \Big|_{z_2^b} \\ &= \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] \frac{1}{z_2^{r_2}} (-1)^{r_1+a} \sum_{s=0}^{r_1+a} e_{r_1+a-s}[X] \frac{1}{z_2^s} e_s[M] \Omega[-z_2 X] \Big|_{z_2^b} \\ &= \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] (-1)^{r_1+a} \sum_{s=0}^{r_1+a} e_{r_1+a-s}[X] e_s[M] (-1)^{r_2+s+b} e_{r_2+s+b}[X] \end{aligned} \quad 1.7$$

where for convenience we have set

$$F^{(r_1)}[X] = F[X + Mu_1] \Big|_{u_1^{r_1}} \quad \text{and} \quad F^{(r_1, r_2)}[X] = F[X + Mu_1 + Mu_2] \Big|_{u_1^{r_1} u_2^{r_2}}$$

We can see from this example that the second calculation of the action of the operator $D_b D_a$ is completely elementary and straight forward. But, in more complex situations, this approach is not conducive to discovery but only to delivering the verification of an identity discovered by other means. On the other hand the calculation of this action carried out in 1.6, in several significant instances, has led to discovery and proof of surprising identities. Nevertheless, we must add that due care must be taken in expressing the rational function, argument of the constant term in 1.6, as an appropriate Laurent series in z_1, z_2 . A systematic way of carrying this out in greater generality has been developed in [30] and [31].

The proof of Theorem I.1 in this section will use the approach illustrated in 1.7. The proof that follows the approach in 1.6, led to the discover of the operators $D_{u,v}$ and their commutator identities. This second proof will be given in section 5.

Recalling that the action of the operator $D_{u,v}$ is defined by setting

$$D_{m,n}F[X] = F[X + [m]_q \frac{M}{z}] \Omega[-z[m]_t X] \Big|_{z^n} \quad 1.8$$

with the convention that $t = 1/q$, we have

Theorem 1.1

If a, b, c, d, m, n are any integers related by the vector identity $(a, b) + (c, d) = (m, n)$ we have for non vanishing a, c, m

$$\frac{1}{M} [D_{c,d}, D_{a,b}] \Big|_{t=1/q} = q^{1+bc} \frac{[a]_q [c]_q}{[m]_q} \frac{1 - q^{da-bc}}{1 - q} D_{m,n} \Big|_{t=1/q}. \quad 1.9$$

Proof

Using the notation and the sequence of steps outlined in 1.7 we get

$$\begin{aligned} D_{a,b}F[X] &= F[X + \frac{M[a]_q}{z_1}] \Omega[-z_1[a]_t X] \Big|_{z_1^b} = \sum_{r_1 \geq 0} F^{(r_1)}[X] \frac{1}{z_1^{r_1}} \Omega[-z_1[a]_t X] \Big|_{z_1^b} \\ &= \sum_{r_1 \geq 0} F^{(r_1)}[X] (-1)^{r_1+b} e_{r_1+b} [[a]_t X] \end{aligned}$$

and consequently

$$\begin{aligned} D_{c,d} D_{a,b} F[X] &= \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] \frac{(-1)^{r_1+b}}{z_2^{r_2}} e_{r_1+b} [[a]_t (X + \frac{M[c]_q}{z_2})] \Omega[-z_2[c]_t X] \Big|_{z_2^d} \\ &= \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] \frac{(-1)^{r_1+b}}{z_2^{r_2}} \sum_{s=0}^{r_1+b} e_{r_1+b-s} [[a]_t X] e_s [[a]_t M[c]_q] \frac{1}{z_2^s} \Omega[-z_2[c]_t X] \Big|_{z_2^d} \\ &= \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] (-1)^{r_1+b} \sum_{s=0}^{r_1+b} e_{r_1+b-s} [[a]_t X] e_s [M[a]_t [c]_q] (-1)^{r_2+d+s} e_{r_2+d+s} [[c]_t X] \end{aligned} \quad 1.10$$

Now we can easy see that

$$M[a]_t [c]_q = \frac{(1-q)(1-1/q)(1-q^{-a})(1-q^c)}{(1-1/q)(1-q)} = -q^{-a}(1-q^a)(1-q^c).$$

Thus

$$e_s [M[a]_t [c]_q] = \frac{(-1)^s}{q^{as}} h_s [(1-q^a)(1-q^c)] = \frac{(-1)^s}{q^{as}} (1-q^a)(1-q^c) \frac{1-q^{sm}}{1-q^m},$$

and 1.10 becomes

$$\begin{aligned} \frac{D_{c,d}D_{a,b}}{(1-q^a)(1-q^c)}F[X] &= \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X](-1)^{r_1+b} \sum_{s=0}^{r_1+b} e_{r_1+b-s}[[a]_t X] \frac{(-1)^s}{q^{as}} \frac{1-q^{sm}}{1-q^m} (-1)^{r_2+d+s} e_{r_2+d+s}[[c]_t X] \\ &= \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X](-1)^{r_1+r_2+n} \sum_{s=0}^{r_1+b} e_{r_1+b-s}[[a]_t X] \frac{q^{-as}-q^{cs}}{1-q^m} e_{r_2+d+s}[[c]_t X] \end{aligned}$$

Or better

$$\frac{(1-q^m)D_{c,d}D_{a,b}}{(1-q^a)(1-q^c)}F[X] = A - B \quad 1.11$$

With

$$A = \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X](-1)^{r_1+r_2+n} \sum_{s=0}^{r_1+b} e_{r_1+b-s}[[a]_t X] q^{-as} e_{r_2+d+s}[[c]_t X]$$

and

$$B = \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X](-1)^{r_1+r_2+n} \sum_{s=0}^{r_1+b} e_{r_1+b-s}[[a]_t X] q^{cs} e_{r_2+d+s}[[c]_t X]$$

Simple manipulation allow us to rewrite A and B in the more convenient forms

$$A = q^{ad} \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] q^{ar_2} (-1)^{r_1+r_2+n} \sum_{s=0}^{r_1+b} e_{r_1+b-s}[[a]_t X] e_{r_2+d+s}[q^{-a}[c]_t X]$$

and

$$B = q^{cb} \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] q^{cr_1} (-1)^{r_1+r_2+n} \sum_{s=0}^{r_1+b} e_{r_1+b-s}[q^{-c}[a]_t X] e_{r_2+d+s}[[c]_t X].$$

Carrying out the interchanges $a \leftrightarrow c$ and $b \leftrightarrow d$ gives

$$\tilde{A} = q^{cb} \sum_{r_1, r_2 \geq 0} F^{(r_2, r_1)}[X] q^{cr_2} (-1)^{r_1+r_2+n} \sum_{s=0}^{r_1+d} e_{r_1+d-s}[[c]_t X] e_{r_2+b+s}[q^{-c}[a]_t X]$$

and

$$\tilde{B} = q^{ad} \sum_{r_1, r_2 \geq 0} F^{(r_2, r_1)}[X] q^{ar_1} (-1)^{r_1+r_2+n} \sum_{s=0}^{r_1+d} e_{r_1+d-s}[q^{-a}[c]_t X] e_{r_2+b+s}[[a]_t X]$$

Thus from 1.11 we derive that

$$\frac{(1-q^m)D_{a,b}D_{c,d}}{(1-q^a)(1-q^c)}F[X] = \tilde{A} - \tilde{B}$$

and consequently

$$\frac{(1-q^m)[D_{c,d}, D_{a,b}]}{(1-q^a)(1-q^c)}F[X] = A - B - (\tilde{A} - \tilde{B}) = (A + \tilde{B}) - (B + \tilde{A}) \quad 1.12$$

Now we may rewrite A by setting $u = r_1 + b - s$ so $0 \leq u \leq r_1 + b$ and $r_2 + d + s = r_1 + r_2 + n - u$ obtaining

$$A = q^{ad} \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] q^{ar_2} (-1)^{r_1+r_2+n} \sum_{u=0}^{r_1+b} e_u[[a]_t X] e_{r_1+r_2+n-u}[q^{-a}[c]_t X]$$

For $\tilde{B}$ we set $u = r_2 + b + s$ so $r_2 + b \leq u \leq r_1 + r_2 + n$ and then make the switch $r_1 \leftrightarrow r_2$

$$\begin{aligned} \tilde{B} &= q^{ad} \sum_{r_1, r_2 \geq 0} F^{(r_2, r_1)}[X] q^{ar_1} (-1)^{r_1+r_2+n} \sum_{u=r_2+b}^{r_1+r_2+n} e_u[[a]_t X] e_{r_1+r_2+n-u}[q^{-a}[c]_t X] \\ &= q^{ad} \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] q^{ar_2} (-1)^{r_1+r_2+n} \sum_{u=r_1+b}^{r_1+r_2+n} e_u[[a]_t X] e_{r_1+r_2+n-u}[q^{-a}[c]_t X] \end{aligned}$$

This gives

$$\begin{aligned}
A + \tilde{B} &= q^{ad} \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] q^{ar_2} (-1)^{r_1+r_2+n} \sum_{u=0}^{r_1+r_2+n} e_u [[a]_t X]] e_{r_1+r_2+n-u} [q^{-a} [c]_t X] \\
&\quad + \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] (-1)^{r_1+r_2+n} e_{r_1+b} [[a]_t X]] e_{r_2+d} [[c]_t X] \\
&= q^{ad} \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] q^{ar_2} (-1)^{r_1+r_2+n} e_{r_1+r_2+n} [[m]_t X] \\
&\quad + \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] (-1)^{r_1+r_2+n} e_{r_1+b} [[a]_t X]] e_{r_2+d} [[c]_t X] \\
&= q^{ad} \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] \frac{q^{ar_2}}{z^{r_1+r_2}} \Omega[-[m]_t X] \Big|_{z^n} \\
&\quad + \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] (-1)^{r_1+r_2+n} e_{r_1+b} [[a]_t X]] e_{r_2+d} [[c]_t X] \\
&= q^{ad} F\left[X + \frac{M[m]_q}{z}\right] \Omega[-[m]_t X] \Big|_{z^n} \\
&\quad + \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] (-1)^{r_1+r_2+n} e_{r_1+b} [[a]_t X]] e_{r_2+d} [[c]_t X].
\end{aligned}$$

With an entirely analogous sequence of steps we obtain the identity

$$\begin{aligned}
B + \tilde{A} &= q^{cb} F\left[X + \frac{M[m]_q}{z}\right] \Omega[-[m]_t X] \Big|_{z^n} \\
&\quad + \sum_{r_1, r_2 \geq 0} F^{(r_1, r_2)}[X] (-1)^{r_1+r_2+n} e_{r_2+d} [[c]_t X]] e_{r_1+b} [[a]_t X]].
\end{aligned}$$

Thus 1.12 finally yields

$$\frac{(1-q^m)[D_{c,d}, D_{a,b}]}{(1-q^a)(1-q^c)} [D_{c,d}, D_{a,b}] F[X] = (q^{ad} - q^{cb}) F\left[X + \frac{M[m]_q}{z}\right] \Omega[-[m]_t X] \Big|_{z^n}.$$

Proving the identity

$$\frac{1}{M} [D_{c,d}, D_{a,b}] F[X] = \frac{q[a]_q [c]_q \frac{q^{cb} - q^{ad}}{1-q}}{[m]_q} F\left[X + \frac{M[m]_q}{z}\right] \Omega[-[m]_t X] \Big|_{z^n},$$

which is just another way of writing 1.9.

In the remainder of this section we will derive a number of immediate consequences of the identity in 1.9. We will state and prove them as a succession of Corollaries, the last two of which are simply restatements of Theorems I.1 and I.2. For convenience we will here and after use the symbol “ $Q_{u,v}^s$ ” as a short hand for “ $Q_{u,v} \Big|_{t=1/q}$ ”. We will start with an auxiliary identity that we will use in various occasions.

Lemma 1.1

For integers a, b, c, d , we have

$$(a-1)(b+1)/2 + (c-1)(d+1)/2 + bc + 1 = (a+c-1)(b+d+1)/2 + (bc - ad + 1)/2.$$

Corollary 1.1

If m and n are relatively prime, then we have

$$Q_{m,n}^s = q^{(n+1)(m-1)/2} D_{m,n} / [m]_q.$$

Proof

We will proceed by induction on m . The base case $m = 1$ is easy: we have $Q_{1,n}^s = D_n \Big|_{t=1/q} = D_{1,n}$, so the corollary holds in this case.

It is best to start by an example. Let us consider $Q_{2,n}^s = \frac{1}{M}[Q_{1,d}^s, Q_{1,b}^s]$, where $Split(2, n) = (1, b) + (1, d)$, with $d - b = 1$. Then by 1.9

$$\begin{aligned} Q_{2,n}^s &= \frac{1}{M}[D_{1,d}, D_{1,b}] = q^{b+1} \frac{[1]_q [1]_q}{[2]_q} \frac{(1-q^{d-b})}{1-q} D_{2,n} \\ &= q^{(m-1)(n+1)/2} \frac{1}{[2]_q} D_{2,n}. \end{aligned}$$

One more example. Let us say $Split(3, n) = (1, b) + (2, d)$ with $d - 2b = 1$. Then again by 1.9

$$\begin{aligned} Q_{3,n}^s &= \frac{1}{M}[Q_{2,d}^s, Q_{1,b}^s] = q^{(d+1)/2} \frac{1}{[2]_q} [D_{2,d}, D_{1,b}] = q^{(d+1)/2} q^{2b+1} \frac{1}{[3]_q} D_{3,n} \\ &= q^{(n+1)} \frac{1}{[3]_q} D_{3,n}. \end{aligned}$$

Assume the corollary holds for smaller m . Now suppose

$$Split(m, n) = (a, b) + (c, d).$$

That is we have $a + c = m, b + d = n, ad - bc = 1$. Then

$$\begin{aligned} Q_{m,n}^s &= \frac{1}{M}[Q_{c,d}^s, Q_{a,b}^s] \\ &= q^{(c-1)(d+1)/2 + (a-1)(b+1)/2} \frac{1}{[a]_q [c]_q} \frac{1}{M}[D_{c,d}, D_{a,b}] \\ &= q^{(c-1)(d+1)/2 + (a-1)(b+1)/2} \cdot q^{bc+1} \frac{1}{[a+c]_q} \frac{1-q^{ad-bc}}{1-q} D_{a+c, b+d} \\ &= q^{(c-1)(d+1)/2 + (a-1)(b+1)/2 + bc+1} \frac{1}{[m]_q} D_{m,n} \\ &= q^{(m-1)(n+1)/2} \frac{1}{[m]_q} D_{m,n}. \end{aligned}$$

This completes the induction and the proof.

Corollary 1.2

For any coprime pair (m, n) we have

$$q^{(m-1)(n-1)/2} Q_{m,n}^s (-1)^n = \frac{1}{[m]_q} e_n[X[m]_q].$$

Proof

$$\begin{aligned} Q_{m,n}^s (-1)^n &= q^{(m-1)(n+1)/2} \frac{1}{[m]_q} D_{m,n} (-1)^n \\ &= q^{(m-1)(n+1)/2} \frac{1}{[m]_q} (-1)^n \Omega - z X[m]_t z^{-n} \Big|_{z^0} \\ &= q^{(m-1)(n+1)/2} \frac{1}{[m]_q} e_n[X[m]_t] \\ &= q^{(m-1)(n+1)/2} q^{-(m-1)n} \frac{1}{[m]_q} e_n[X[m]_q] \\ &= q^{-(m-1)(n-1)/2} \frac{1}{[m]_q} e_n[X[m]_q]. \end{aligned}$$

Corollary 1.3

For any coprime pair (m, n) and $k \geq 1$ we have

$$Q_{km, kn}^s = q^{(km-1)(kn+1)/2 - (k-1)/2} \frac{[k]_q}{[km]_q} D_{km, kn}.$$

Proof

Suppose $Split(m, n) = (a, b) + (c, d)$. That is $a + c = m, b + d = n, ad - bc = 1$. It should be clear that a, b are also relatively prime, we can choose $Split(km, kn) = (a, b) + ((k-1)a + kc, (k-1)b + kd)$, so that it is easily obtained by linear algebra that

$$\det \begin{pmatrix} a & (k-1)a + kc \\ b & (k-1)b + kd \end{pmatrix} = k, \quad \det \begin{pmatrix} a+c & (k-1)a + kc \\ b+d & (k-1)b + kd \end{pmatrix} = 1,$$

Thus $c' = (k-1)a + kc$ and $d' = (k-1)b + kd$ are relatively prime. We have

$$\begin{aligned} Q_{km, kn}^s &= \frac{1}{M} [Q_{(k-1)a+kc, (k-1)b+kd}^s, Q_{a, b}^s] \\ &= q^{(c'-1)(d'+1)/2+(a-1)(b+1)/2} \frac{1}{[a]_q [c']_q} \frac{1}{M} [D_{c', d'}, D_{a, b}] \\ &= q^{(c'-1)(d'+1)/2+(a-1)(b+1)/2} \cdot q^{bc'+1} \frac{1}{[a+c']_q} \frac{1-q^{ad'-bc'}}{1-q} D_{a+c', b+d'} \\ &= q^{(c'-1)(d'+1)/2+(a-1)(b+1)/2+bc'+1} \frac{[k]_q}{[km]_q} D_{km, kn} \\ &= q^{(km-1)(kn+1)/2-(k-1)/2} \frac{[k]_q}{[km]_q} D_{km, kn}. \end{aligned}$$

Corollary 1.4

For any coprime pair (m, n) and $k \geq 1$ we have

$$Q_{km, kn}^s (-1)^{kn} = q^{-(km-1)(kn-1)/2-(k-1)/2} \frac{[k]_q}{[km]_q} e_n [X[km]].$$

Proof

We have

$$\begin{aligned} Q_{km, kn}^s (-1)^{kn} &= q^{(km-1)(kn+1)/2-(k-1)/2} \frac{[k]_q}{[km]_q} D_{km, kn} (-1)^{kn} \\ &= q^{(km-1)(kn+1)/2-(k-1)/2} \frac{[k]_q}{[km]_q} D_{km, kn} (-1)^{kn} (-1)^{kn} \Omega_{-z} X[km]_t \Big|_{z^{kn}} \\ &= q^{(km-1)(kn+1)/2-(k-1)/2} \frac{[k]_q}{[km]_q} e_n [X[km]_t] \\ &= q^{(km-1)(kn+1)/2-(k-1)/2} q^{-(km-1)n} \frac{[k]_q}{[km]_q} e_n [X[km]_q] \\ &= q^{-(km-1)(kn-1)/2-(k-1)/2} \frac{[k]_q}{[km]_q} e_n [X[km]_q]. \end{aligned}$$

2. Polynomiality and positivity.

The main goal of this section is to prove that the quotient

$$\frac{[k]_q}{[km]_q} e_{kn} [X[km]_q] \quad 2.1$$

is a Schur positive symmetric polynomial. This will be obtained by combining the next four auxiliary propositions.

The first fact on which the proof is based is the following classical result.

Proposition 2.1

For any $n \geq 1$ we have to matrices $\|c_{\lambda,\rho}\|_{\lambda,\rho \vdash n}$ and $\|d_{\lambda,\rho}\|_{\lambda,\rho \vdash n}$ such that

$$a) \quad s_\lambda = \sum_{\rho \vdash n} c_{\lambda,\rho} p_\rho \quad \text{and} \quad b) \quad p_\rho = \sum_{\lambda \vdash n} d_{\lambda,\rho} s_\lambda \quad 2.2$$

Proof

Frobenius proved that 2.2 a) and b) hold with

$$a) \quad c_{\lambda,\rho} = \chi_\rho^\lambda / z_\rho \quad \text{and} \quad d) \quad d_{\lambda,\rho} = \chi_\rho^\lambda \quad 2.3$$

with χ_ρ^λ the value of Young's irreducible S_n character indexed by λ at the conjugacy class indexed by ρ and for $\rho = 1^{\alpha_1} 2^{\alpha_2} \dots n^{\alpha_n}$ we have

$$z_\rho = 1^{\alpha_1} 2^{\alpha_2} \dots n^{\alpha_n} \alpha_1! \alpha_2! \dots \alpha_n!$$

This given, our polynomiality result can be stated as follows

Proposition 2.2

If $(m, n) = 1$ and $k \geq 1$ then for all $\lambda \vdash kn$ we have

$$\frac{[k]_q}{[km]_q} s_\lambda [[km]_q] \in \mathbb{Q}[q] \quad 2.4$$

where for any integer $s \geq 0$ we set $[s]_q = 1 + q + \dots + q^{s-1}$

Proof

Note that to show 2.4 we need only show that every root of

$$1 + q + q^2 + \dots + q^{km-1} = 0 \quad 2.5$$

is a root of the polynomial

$$[k]_q s_\lambda [[km]_q].$$

To show this we use 2.2 (for $n \rightarrow kn$) and write for ζ a root of 2.5

$$[k]_\zeta s_\lambda [[km]_\zeta] = \sum_{\rho \vdash kn} c_{\lambda,\rho} [k]_\zeta p_\rho [[km]_\zeta] \quad 2.6$$

Thus if the left hand side of 2.6 does not vanish we will necessarily have a $\rho \vdash kn$ such that

$$[k]_\zeta p_\rho [[km]_\zeta] \neq 0 \quad 2.7$$

in particular if for some r we have $\rho_r = i$ then

$$1 + \zeta^i + \zeta^{2i} + \dots + \zeta^{(km-1)i} = p_i [[km]_\zeta] \neq 0 \quad 2.8$$

and since

$$(1 - q^i)(1 + q^i + q^{2i} + \cdots + q^{(km-1)i}) = 1 - q^{(km)i}$$

from 2.8, and the fact that ζ is a root of 2.5, we derive that

$$a) \quad \zeta^i = 1 \quad \& \quad b) \quad \zeta \neq 1 \quad 2.9$$

Now note that since the ρ in 2.7 may be written in the form $\rho = \prod_{i=1}^{kn} i^{\alpha_i}$ with $\sum_{i=1}^{kn} i\alpha_i = kn$ then we must also have

$$\zeta^{kn} = \prod_{i=1}^{kn} (\zeta^i)^{\alpha_i} = 1$$

Now the assumed primality of the pair m, n gives that $k = \gcd(km, kn)$ and this combined with the fact that ζ is a root of 2.5 forces

$$\zeta^k = 1$$

but then b) of 2.9 yields

$$1 + \zeta + \zeta^2 + \cdots + \zeta^{k-1} = 0$$

which is in plain contradiction with 2.7. This contradiction forces every root of 2.5 to be a root of

$$[k]_q s_\lambda [[km]_q] = 0$$

as desired.

The next device we use is the following well known fact

Proposition 2.3

Any principal evaluation of a Schur Function is unimodal. More precisely for any $\lambda \vdash n$ and any $m > 1$ the polynomial

$$s_\lambda [[m]_q] \quad 2.10$$

is unimodal.

Proof

This is exercise 4 page 137 of Macdonald book [25]. Since the solution in [25] is barely sketched, for sake of completeness we carry out Macdonald's exercise in full detail. Macdonald considers the evaluation

$$s_\lambda \left[\frac{x_1^m - x_2^m}{x_1 - x_2} \right] = s_\lambda [s_{m-1}[x_1 + x_2]] \quad 2.11$$

as a character of $GL_2[\mathbf{C}]$. Using this he derives that for some weakly positive integer constants c_{r_1, r_2} we must have the expansion

$$s_\lambda [s_{m-1}[x_1 + x_2]] = \sum_{\substack{r_1 + r_2 = d \\ r_1 \geq r_2}} c_{r_1, r_2} s_{[r_1, r_2]} [x_1 + x_2] \quad 2.12$$

where for convenience we have set

$$d = (m-1)n \quad 2.13$$

Notice that we have

$$\begin{aligned} s_{[r_1, r_2]} [x_1 + x_2] &= x_1^{r_1} x_2^{r_2} + x_1^{r_1-1} x_2^{r_2+1} + \cdots + x_1^{r_2+1} x_2^{r_1-1} + x_1^{r_2} x_2^{r_1} \\ &= x_2^d \left(\left(\frac{x_1}{x_2} \right)^{r_1} + \left(\frac{x_1}{x_2} \right)^{r_1-1} + \cdots + \left(\frac{x_1}{x_2} \right)^{r_2} \right) = x_2^d \sum_{s=r_2}^{r_1} q^s \end{aligned}$$

where for convenience we have set $\frac{x_1}{x_2} = q$. Thus with a slight change of notation we may rewrite 2.12 in the form

$$s_\lambda[s_{m-1}[x_1 + x_2]] = x_2^d \sum_{r_2=0}^{\lfloor d/2 \rfloor} c_{r_2} \sum_{s=0}^d q^s \chi(r_2 \leq s \leq d - r_2) = x_2^d \sum_{s=0}^d q^s \sum_{r_2=0}^{s \wedge (d-s)} c_{r_2} \quad 2.14$$

from which the unimodality assertion immediately follows by setting $x_2 = 1$ and $x_1 = q$.

Our positivity result is a consequence of the following simple but powerful fact

Proposition 2.4

Let $g(q) = b_0 + b_1 q + \cdots + b_r q^r$ and assume that, for $d = r + s$, the polynomial

$$f(q) = (1 + q + \cdots + q^s)g(q) = \sum_{l=0}^d c_l q^l \quad 2.15$$

is unimodal with peak at p and non-negative coefficients. Then $g(q)$ also has non-negative coefficients.

Proof

We proceed by contradiction. Suppose some of the coefficients of $g(b)$ are negative. Since $c_0 = c_d \geq 0$ let b_i and b_j (with $0 < i \leq j < d$) be the leftmost and rightmost negative coefficients of $g(q)$ respectively. Now if $i \leq p$ then

$$c_i = \sum_{u=0 \vee (i-s)}^i b_u \quad \& \quad c_{i-1} = \sum_{u=0 \vee (i-1-s)}^{i-1} b_u$$

This gives

$$c_i - c_{i-1} = b_i - \chi(i-1-s > 0) b_{i-1-s} < 0$$

a contradiction!

If $r-j \leq d-p$ then do the same argument for $\tilde{f}(q) = q^d f(1/q)$ and $\tilde{g}(q) = q^r g(1/q)$. So we are left with $i > p$ and $r-j > d-p$. But that cannot happen since it implies that $i > s + j > j$.

As a corollary we obtain our desired goal.

Theorem 2.1

For any coprime pair (m, n) and any $k \geq 1$ we have that the symmetric function

$$\frac{[k]_q}{[km]_q} e_{kn} [X[km]_q] \quad 2.16$$

is a Schur positive symmetric polynomial

Proof

The Cauchy formula gives

$$\frac{[k]_q}{[km]_q} e_{kn} [X[km]_q] = \sum_{\lambda \vdash kn} s_{\lambda'} [X] \frac{[k]_q s_{\lambda} [[km]_q]}{[km]_q}$$

Now we have proved that $[k]_q s_{\lambda} [[km]_q]$ is divisible by $[km]_q$, we have also proved that $s_{\lambda} [[km]_q]$ is palindromic unimodal. Thus it follows from this that also $[k]_q s_{\lambda} [[km]_q]$ is palindromic unimodal, we can then apply Proposition 2.4 with $f(q) = [k]_q s_{\lambda} [[km]_q]$ and

$$g(q) = \frac{[k]_q s_{\lambda} [[km]_q]}{[km]_q}$$

and conclude that $g(q) \in \mathbf{N}[q]$, proving the Schur positivity of the polynomial in 2.16.

Remark 2.1

The polynomiality of the symmetric function in 2.16 for the special case $k = 1$ was first proved by Mark Haiman in [22]. The question arose from the discovery in [10] that in the case $m = n + 1$ 2.16 (for $k = 1$) is the Frobenius characteristic of an appropriate single grading of the Diagonal Harmonic Module of S_n . This also prompted Haiman to seek for some reason that justified this polynomial to be Schur positive. With remarkable foresight Haiman investigates the more general (m, n) case and provided a mechanism for proving that the symmetric function

$$\frac{e_n[X[m]_q]}{[m]_q} \tag{2.17}$$

is a polynomial and Schur positive if and only if (m, n) is a coprime pair. However, the Schur positivity was shown in [22] by constructing a quotient

$$\mathbb{Q}[x_1, x_2, \dots, x_n] / (e_1, f_2, \dots, f_n)$$

with Frobenius characteristic the polynomial in 2.17, where e_1 the ordinary elementary symmetric function and $f_1, f_2, \dots, f_n$ a sequence of polynomials satisfying the following properties

- (1) *each f_i is homogeneous of degree m*
- (2) *they satisfy the identities $\sigma f_i = f_{\sigma_i}$ (for $1 \leq i \leq n$ and all $\sigma \in S_n$)*
- (3) $f_1 + f_2 + \dots + f_n = 0$
- (4) $e_1, f_2, \dots, f_n$ are a regular sequence.

A sequence f_i satisfying (1),(2),(3),(4) was constructed by Hanspeter Kraft a few years later but never published. More recently a very natural example of such a sequence was discovered by Dunkl in [6] and used later by Gorsky in his work [16] on torus knots invariants. It follows from this and Mark Haiman result the Schur positivity of the polynomial in 2.17 in the coprime case.

The challenge now is to construct an equally natural quotient with Frobenius characteristic the polynomial in 2.16.

3. A Parking Function setting for our Frobenius Characteristics

The main goal of this section is a proof of Theorem I.4. To carry this out we need to briefly review the statement of the Rational Compositional Shuffle Conjecture. We will start by introducing the symmetric function tools that are used in its formulation.

The basic ingredient here is the identity

$$Q_{0,k} = \frac{qt}{qt-1} \underline{h}_k[X(1/qt - 1)] \quad (\text{for all integers } k \geq 1). \quad 3.1$$

From this it follows that a basis for the subspace of operators of bi-degrees $(0, n)$ for $n \geq 1$ is given by the collection

$$\{Q_{0,\lambda}\}_\lambda = \left\{ \prod_{i=1}^{l(\lambda)} Q_{0,\lambda_i} \right\}_\lambda \quad 3.2$$

This fact, (see [3] for an elementary treatment), can be used to construct an operator of bi-degree (km, kn) for any coprime pair (m, n) , any integer $k \geq 1$ and any given homogeneous symmetric function $F[X]$ of degree k , by the following two steps.

(1) *Compute the expansion*

$$F = \sum_{\lambda \vdash k} c_\lambda \prod_{i=1}^{l(\lambda)} Q_{0,\lambda_i}, \quad 3.3$$


(2) *and then set*

$$\mathbf{F}_{km,kn} = \sum_{\lambda \vdash k} c_\lambda \prod_{i=1}^{l(\lambda)} Q_{\lambda_i m, \lambda_i n}. \quad 3.4$$

The commutativity of the operators Q_{u_1, v_1} and Q_{u_2, v_2} with (u_1, v_1) and (u_2, v_2) collinear vectors assures that 3.4 well defines the operator $\mathbf{F}_{km,kn}$. In [3] a variety of Shuffle conjectures were formulated based on the above construction and various choices of the symmetric function $F[x]$. The simplest one corresponds to choosing $F = e_k$ (the k^{th} elementary symmetric function). The corresponding operator which we denote $\mathbf{e}_{km,kn}$ has a truly remarkable connection to the Theory of Parking Functions.

A single example will suffice to get across this connection. We have displayed here on the right a 12×20 lattice rectangle with a path that proceeds by North and East unit steps, always remaining weakly above the diagonal $(0,0) \rightarrow (12,20)$. A Parking Function in the 12×20 lattice rectangle is obtained by labeling the cells immediately to the right of the north steps of such a path by the integers $1, 2, \dots, 20$ (referred to as *cars*) in a column increasing manner. This given, one of the conjectures formulated in [3] may be expressed as the identity

$$\mathbf{e}_{km,kn}(-1)^{k(n+1)} = \sum_{PF \in \mathcal{PF}_{km,kn}} t^{\text{area}(PF)} q^{\text{dinv}(PF)} F_{\text{pides}(PF)} \quad 3.5$$



where the sum is over all Parking Functions in the $km \times kn$ lattice rectangle, and the Parking Function statistics occurring in 3.5 are as will be defined shortly. Here the symbol “ $F_{\text{pides}(PF)}$ ” stands for the Gessel’s [15] fundamental quasi-symmetric function indexed by the composition “ $\text{pides}(PF)$ ”.

Now the Extended Compositional Shuffle Conjecture states a refined version of 3.5 where the symmetric function F in the above construction of the operator $\mathbf{F}_{km,kn}$, is chosen to assure that the sum in 3.5 is restricted to be over an appropriately selected subfamily of Parking Functions in the $km \times kn$ lattice rectangle. These special choices of F are obtained by means of the modified Hall-Littlewood operators C_a whose action on a symmetric function $F[X]$ is defined by setting

$$C_a P[X] = \left(-\frac{1}{q}\right)^{a-1} P\left[X - \frac{1-1/q}{z}\right] \sum_{m \geq 0} z^m h_m[X] \Big|_{z^a} \quad 3.6$$

Again, a single example will suffice to illustrate our choices. Suppose that we want to restrict the sum in 3.5 to be carried out only over the Parking Functions whose supporting Dyck path hits the diagonal precisely at the first and third and fourth possible places, as the path depicted in the above display. To achieve this we simply choose the symmetric function $F = C_1 C_2 C_1 1$. More generally, given a composition $p = (p_1, p_2, \dots, p_l)$ of the integer k , let us denote by $\mathbf{C}_{km, kn}^{(p)}$ the operator obtained by choosing $F = C_{p_1} C_{p_2} \cdots C_{p_l} 1$ in the above construction. This given, the Extended Compositional Shuffle Conjecture states that

$$\mathbf{C}_{km, kn}^{(p)} (-1)^{k(n+1)} = \sum_{PF \in \mathcal{PF}_{km, kn}(p)} t^{\text{area}(PF)} q^{\text{area}(PF)} F_{\text{pides}(PF)} \quad 3.7$$

where the sum is over Parking Functions whose path hits the diagonal precisely in l of the k possible places, as prescribed by the parts of the composition $p = (p_1, p_2, \dots, p_l)$. The reason 3.7 refines 3.6 is due to the remarkable identity

$$e_k = \sum_{p \models k} C_{p_1} C_{p_2} \cdots C_{p_l(p)} 1.$$

Here the sum is over all compositions of k

Keeping all this in mind, we are now in a position to show that the identity in I.17 is one of the many consequences of the identity in 3.7. More precisely we will show that Theorem I.4 is a corollary of the following stronger result.

Theorem 3.1

The validity of 3.7 for any coprime pair (m, n) , $k \geq 1$ and any composition $p \models k$ implies the identity

$$Q_{km, kn} (-1)^{kn} = \sum_{PF \in \mathcal{PF}_{km, kn}} t^{\text{area}(PF) - \text{ret}(PF) + 1} [\text{ret}(PF)]_t q^{\text{dinv}(PF)} F_{\text{pides}(PF)} \quad 3.8$$

where all the Parking Function statistics are as in the Extended Shuffle Conjectures. The “ $\text{ret}(PF)$ ” statistic gives the smallest positive i such that the supporting path of PF goes through the point (im, in) .

Proof

For brevity we will start with the following identity, valid for any integer $1 \leq d \leq k$ and $p \models k - d$

$$\sum_{p \models k-d} C_d C_{p_1} C_{p_2} \cdots C_{p_l(p)} 1 = \left(-\frac{1}{q}\right)^{d-1} s_{d, 1^{k-d}} - \left(-\frac{1}{q}\right)^d s_{1+d, 1^{k-d-1}} \quad 3.9$$

(see [21] for a proof). Note next that it follows from 3.9 that

$$\sum_{d=a}^k \sum_{p \models k-d} C_d C_p 1 = \sum_{d=a}^k \left(-\frac{1}{q}\right)^{d-1} s_{d, 1^{k-d}} - \sum_{d=a+1}^k \left(-\frac{1}{q}\right)^{d-1} s_{d, 1^{k-d}} = \left(-\frac{1}{q}\right)^{a-1} s_{a, 1^{k-a}} \quad 3.10$$

Our next step is to rewrite 3.1 in a more suitable form. Now a use of the Cauchy identity gives

$$\begin{aligned} Q_{0,k} &= \frac{(qt)^{1-k}}{qt-1} h_k[X(1-qt)] = \frac{(qt)^{1-k}}{qt-1} \sum_{\mu \vdash k} s_\mu[X] s_\mu[(1-qt)] \\ &= \frac{(qt)^{1-k}}{qt-1} \sum_{r=0}^{k-1} s_{k-r, 1^r}[X] (-qt)^r (1-qt) = -(qt)^{1-k} \sum_{r=0}^{k-1} s_{k-r, 1^r}[X] (-qt)^r \\ &= (-1)^k \sum_{a=1}^k s_{a, 1^{k-a}}[X] (-qt)^{1-a}, \end{aligned} \quad 3.11$$

where for the third equality we use the following, easily verified, special evaluation of a Schur function

$$s_\mu[1-m] = \begin{cases} (-m)^a (1-m) & \text{if } \mu = k-r, 1^r \text{ for } 0 \leq r \leq k-1 \\ 0 & \text{otherwise} \end{cases}, \quad 3.12$$

valid for any monomial m .

Hence by combining 3.10 and 3.11 we derive that

$$\begin{aligned} (-1)^k Q_{0,k} &= \sum_{a=1}^k (-qt)^{1-a} s_{a,1^{k-a}} = \sum_{a=1}^k t^{1-a} \sum_{d=1}^k \chi(d \geq a) \sum_{p \models k-d} C_d C_p 1 \\ &= \sum_{d=1}^k \sum_{p \models k-d} C_d C_p 1 \sum_{a=1}^d (1/t)^{a-1} = \sum_{d=1}^k \sum_{p \models k-d} [d]_{1/t} C_d C_p 1 \end{aligned} \quad 3.13$$

This given, the particular case $F = Q_{0,k}$ of the above construction gives that for any coprime pair (m, n) we have

$$\mathbf{Q}_{\mathbf{0}, \mathbf{k}_{km, kn}} = Q_{km, kn} \quad 3.14$$

Likewise by choosing $F = C_d C_p 1$ for $p \models k - d$ we obtain the operator $\mathbf{C}_{\mathbf{d}, \mathbf{p}} \mathbf{1}_{km, kn}$ which, by the Rational Shuffle Compositional conjecture, satisfies

$$\mathbf{C}_{\mathbf{d}, \mathbf{p}} \mathbf{1}_{km, kn} (-1)^{k(n+1)} = \sum_{PF \in \mathcal{PF}_{km, kn}(d)} t^{\text{area}(PF)} q^{\text{div}(PF)} F_{\text{pides}(PF)} \quad 3.15$$

where the sum is over Parking Functions in the $km \times kn$ rectangle whose supporting path returns to the diagonal for the first time in row dn . Thus combining 3.15 with 3.14 and 3.13 we obtain that

$$Q_{km, kn} (-1)^{kn} = \sum_{d=1}^k \sum_{p \models k-d} [d]_{1/t} \sum_{PF \in \mathcal{PF}_{km, kn}(d)} t^{\text{area}(PF)} q^{\text{div}(PF)} F_{\text{pides}(PF)}$$

which is only another way of writing 3.8.

We can now finally give our

Proof of Theorem I.4

We will show here the identity in I.17, written in the form

$$\frac{[k]_q}{[km]_q} e_{kn} [X[km]_q] = \sum_{PF \in \mathcal{PF}_{km, kn}} q^{\text{coarea}(PF) + \text{div}(PF)} [\text{ret}(PF)]_q F_{\text{pides}(PF)} [X] \quad 3.16$$

To this end, notice that setting $t = 1/q$ in 3.8 gives

$$Q_{km, kn} (-1)^{kn} \Big|_{t=1/q} = \sum_{PF \in \mathcal{PF}_{km, kn}} q^{-\text{area}(PF) + \text{ret}(PF) - 1} [\text{ret}(PF)]_q q^{-(\text{ret}(PF) - 1)} q^{\text{div}(PF)} F_{\text{pides}(PF)} \quad 3.17$$


But we can now use I.15 which can be rewritten in the form

$$\frac{[k]_q}{[km]_q} e_{kn} [X[km]_q] = q^{(km-1)(kn-1)/2 + (k-1)/2} Q_{km, kn} (-1)^{kn} \Big|_{t=1/q} \quad 3.18$$

This given, a comparison of the combination of 3.18 and 3.17 with 3.16 shows that we need only to show the equality

$$\text{coarea}(PF) + \text{area}(PF) = (km - 1)(kn - 1)/2 + (k - 1)/2 \quad 3.19$$

This can be easily justified by the following geometric argument. We have depicted in the display above the $km \times kn$ lattice rectangle for $k = 4$ and $(m, n) = (5, 7)$. Now by the definition $\text{area}(PF)$ gives the number of lattice cells below the supporting path of PF and weakly above the diagonal $(0, 0) \rightarrow (km, kn)$ and $\text{coarea}(PF)$ gives the number of lattice cells cells above the path. Thus to show 3.19 we need only verify that the right hand side gives the number of lattice cells weakly above the diagonal $(0, 0) \rightarrow (km, kn)$. Now it



is easy to see from the display that the number of lattice cells cut by the diagonal in any one of the diagonal 4×7 blocks is by one short of $4 + 7$. This implies that the number of uncut lattice cells above the diagonal within each diagonal block is none other than $(mn - m - n + 1)/2 = (m - 1)(n - 1)/2$. Moreover, the total number of lattice cells within the upper non-diagonal blocks is $mn \times \binom{k}{2}$. Thus 3.19 is none other than a consequence of the equality


$$k \times (m - 1)(n - 1)/2 + mn \times \binom{k}{2} = (km - 1)(kn - 1)/2 + (k - 1)/2$$

This completes our proof of 3.16.

In the remainder of this section we will present the latest version of the Parking Function statistics that occur in the various formulations of the rational Compositional Shuffle Conjecture.

Let (m, n) be coprime pair of positive integers and let $k \geq 1$. Recall that a km, kn -Dyck path is sequence of north and east steps in the $km \times kn$ lattice rectangle which starts in the southwest corner, ends in the northeast corner, and stays weakly above the main diagonal $y = \frac{n}{m}x$. A km, kn -parking function is a km, kn -Dyck path with labels $\{1, 2, \dots, kn\}$, known as cars, adjacent to north steps and increasing within each column. For example, see the 6, 9-parking function below. Let $\mathcal{PF}_{km, kn}$ denote the set of all km, kn -parking functions.

The original Rational Shuffle Conjecture of [19] states that for coprime m, n , we can express $Q_{m, n}(-1)^n$ as a weighted sum of m, n -parking functions. This enumeration involves the statistics $area(PF)$, $dinv(PF)$, and the word $\sigma(PF)$. The simplest of these is $area(PF)$, which is the number of full cells between the path and the main diagonal $y = \frac{n}{m}x$. In the adjacent example, the area is 5 and the corresponding cells are shaded.



The $dinv$ and word statistics both make use of a rank function. Let $rank(x, y) = kmy - knx + \lfloor \frac{x}{m} \rfloor$. This causes the points weakly above the diagonal to have distinct nonnegative ranks, with points further from the main diagonal having higher rank. This way, the rank function generalizes the notion of diagonals from classical Parking Function theory [20]. Let the rank of a car be the rank of the southwest corner of that car's cell. Then the word, $\sigma(PF)$, is just the permutation of $\{1, 2, \dots, kn\}$ obtained by listing the cars from highest to lowest rank. In the example above, the ranks of cars 1 through 9 are 9, 0, 6, 13, 2, 8, 14, 15, and 12, respectively. Hence the word of that parking function is $\sigma(PF) = 874916352$.

We set

$$tdinv(PF) = \sum_{\text{cars } i < j} \chi(rank(i) < rank(j) < rank(i) + km). \quad 3.20$$

Here $tdinv$ is short for "temporary $dinv$ " because we will modify this statistic to obtain $dinv(PF)$. In the example above, $tdinv(PF) = 9$ because the inequalities in 3.20 are satisfied for the pairs $(1, 4)$, $(1, 7)$, $(1, 9)$, $(2, 5)$, $(3, 6)$, $(4, 7)$, $(4, 8)$, $(6, 9)$ and $(7, 8)$.

The original formulation of Hikita [24] as modified by Gorsky-Mazin [17], [18] expressed the $dinv$ statistic as a combination of $tdinv$ and two other statistics. However, Hicks and Leven [23] showed that this can be simplified as follows. Let $\lambda(PF)$ be partitions whose english Ferrers diagram is formed by the cells above PF . In the example above, $\lambda(PF) = (4, 4, 4, 2, 1, 1)$. This given, for an m, n -Parking Function we set

$$dinv(PF) = \begin{cases} tdinv(PF) - \# \left\{ c \in \lambda(PF) : \frac{arm(c)}{leg(c)} \leq \frac{m}{n} < \frac{arm(c)+1}{leg(c)+1} \right\} & \text{if } m < n \\ tdinv(PF) + \# \left\{ c \in \lambda(PF) : \frac{arm(c)}{leg(c)} > \frac{m}{n} \geq \frac{arm(c)+1}{leg(c)+1} \right\} & \text{if } m > n. \end{cases} \quad 3.21$$

Here we must use the conventions $\frac{0}{0} = 0$ and $\frac{x}{0} = \infty$ when $x \neq 0$.

In the example above, we have $m < n$, thus $\text{dinv}(PF) = t\text{dinv}(PF) - 4 = 5$, since there are 3 cells with $\text{arm} = 0$ and $\text{leg} = 0$ and one cell with $\text{arm} = 2$ and $\text{leg} = 3$.

We now have all the ingredients that occur in any of the rational Shuffle Conjectures including the Compositional ones in [3]. In particular, 3.5 conjectures the equality

$$\mathbf{e}_{km, kn}(-1)^{k(n+1)} = \sum_{PF \in \mathcal{PF}_{km, kn}} t^{\text{area}(PF)} q^{\text{dinv}(PF)} F_{\text{pides}(PF)} \quad 3.22$$

where $\text{pides}(PF)$ is the composition that gives the descent set of the inverse of the permutation $\sigma(PF)$ defined above, and $F_{\text{pides}(PF)}$ is the Gessel [15] fundamental quasi-symmetric function indexed by the composition $\text{pides}(PF)$. Here we use the convention that for a composition $p \models u$ if $S(p)$, is the subset of $\{1, 2, \dots, u-1\}$ that corresponds to p , then we set

$$F_p(x_1, x_2, \dots, x_v) = \sum_{\substack{1 \leq a_1 \leq a_2 \leq \dots \leq a_v \leq v \\ i \in S(p) \rightarrow a_i < a_{i+1}}} x_{a_1} x_{a_2} \cdots x_{a_v}$$

The same conventions apply to the statistics occurring in 3.8, namely our conjecture that

$$Q_{km, kn}(-1)^{kn} = \sum_{PF \in \mathcal{PF}_{km, kn}} t^{\text{area}(PF) - \text{ret}(PF) + 1} [\text{ret}(PF)]_t q^{\text{dinv}(PF)} F_{\text{pides}(PF)} \quad 3.23$$

Remark 3.1

We must mention that the conjectured equality in 3.16 has a specialization that extends the equality in I.1 to the non coprime case. More precisely, we have

$$\sum_{D \in \mathcal{D}_{km, kn}} [\text{ret}(PF)]_q q^{\text{coarea}(D) + \text{dinv}(D)} = \frac{[k]_q}{[km]_q} \begin{bmatrix} kn + km - 1 \\ kn \end{bmatrix}_q \quad 3.24$$

In fact, scalar multiplication of both sides of 3.16 by $e_{kn}[X]$ gives

$$\begin{aligned} \sum_{D \in \mathcal{D}_{km, kn}} [\text{ret}(PF)]_q q^{\text{coarea}(D) + \text{dinv}(D)} &= \frac{[k]_q}{[km]_q} \langle e_{kn}[X[km]_q], e_{kn}[X] \rangle \\ &= \frac{[k]_q}{[km]_q} \langle h_{kn}[X[km]_q], h_{kn}[X] \rangle \end{aligned}$$

and 3.24 then follows from the identities

$$\langle h_{kn}[X[km]_q], h_{kn}[X] \rangle = h_{kn}[[km]_q] = \begin{bmatrix} kn + km - 1 \\ kn \end{bmatrix}_q$$


We should mention that, for Dyck paths, the dinv statistic is obtained by counting the number of cells c of the english partition above the path (see adjacent figure) whose arm and leg satisfy the inequalities

$$\frac{\text{arm}(c)}{\text{leg}(c) + 1} \leq \frac{m}{n} < \frac{\text{arm}(c) + 1}{\text{leg}(c)}$$

In the example above we have placed a green square in each of the cells that contribute to the dinv .

Remark 3.2

This should complete our presentation of the combinatorial side of the rational Shuffle Conjectures except for two important observations. Firstly we should notice that 3.16 and I.17 differ in that I.16 has “ $s_{\text{pides}(PF)}$ ” replacing “ $F_{\text{pides}(PF)}$ ” in 3.16. We stated I.16 and various analogous identities in the



introduction in this manner, since this makes them easier to verify on a computer. In fact, the validity of this replacement, is one of the surprising consequences of a result of Egge, Loehr and Warrington [7] concerning Gessel fundamental expansions of symmetric functions.

The second observation results from a direct comparison of 3.22 and 3.23. Notice that we have $1 \leq \text{ret}(PF) \leq k$ since the path must end at the point (km, kn) . Furthermore, $\text{area}(PF) \geq \text{ret}(PF) - 1$. This is because each time PF fails to touch the point (im, in) , a cell must fall between the path and the main diagonal. Therefore all the powers of t appearing in 3.23 when $\text{ret}(PF) > 1$ are non-negative. It follows from this that the difference of the right and sides of 3.23 and 3.22 can be shown to be positive linear combination of LLT polynomials which have in turn been shown to be Schur positive. So another strong evidence supporting the validity of these conjectures is that computer data confirms the Schur positivity of the difference of the left hand sides of 3.23 and 3.22.

4. The action of the operators $D_{u,v}$ on the basis $\{s_\mu[\frac{X}{1-q}]\}_\mu$

Our main goals in this section are the proofs of Theorems I.5 and I.6. To carry this out we need auxiliary notation and some preliminary identities which may initially appear only remotely connected with these goals.

Our basic tool here is a new variant of the Macdonald operator “ D_n^1 ” of [25]. We will denote it “ R_v ”. Its action on a symmetric polynomial $P[X_n]$ is obtained by setting

$$R_v P[X_n] = \frac{1}{\Delta[X_n]} \sum_{i=1}^n T_{x_i}^q x_i^v \Delta[X_n] P[X_n], \quad 4.1$$

where “ $T_{x_i}^q$ ” is the linear operator which carries out the substitution $x_i \rightarrow qx_i$. Here $\Delta[X_n]$ is the Vandermonde determinant

$$\Delta[X_n] = \prod_{1 \leq i < j \leq n} (x_i - x_j) = \sum_{\sigma \in S_n} \epsilon(\sigma) \sigma x^\delta, \quad 4.2$$

with $\epsilon(\sigma)$ the sign of σ and $\delta = (n-1, n-2, \dots, 1, 0)$. The actual value of n in both 4.1 and 4.2 is immaterial provided that we choose it greater than v plus the degree of P . The following identity, which in particular shows that R_v preserves symmetry, will play a crucial role.

Proposition 4.1

For any integral vector $\mu = (\mu_1 \geq \mu_2 \geq \dots \geq \mu_n \geq 0)$ we have

$$R_v s_\mu[X_n] = \sum_{i=1}^n q^{v+\mu_i+n-i} s_{\mu+ve_i}[X_n] \quad 4.3$$

with e_i the n dimensional coordinate vector with i^{th} component equal to 1.

Proof

Since by definition

$$s_\mu[X_n] = \frac{1}{\Delta[X_n]} \sum_{\sigma \in S_n} \epsilon(\sigma) \sigma x^{\mu+\delta}$$

we may write

$$\begin{aligned} R_v s_\mu[X_n] &= \frac{1}{\Delta[X_n]} \sum_{i=1}^n T_{x_i}^q x_i^v \sum_{\sigma \in S_n} \epsilon(\sigma) \sigma x^{\mu+\delta} \\ &= \frac{1}{\Delta[X_n]} \sum_{i=1}^n \sum_{\sigma \in S_n} \epsilon(\sigma) T_{x_{\sigma_i}}^q x_{\sigma_i}^v \sigma x^{\mu+\delta} = \frac{1}{\Delta[X_n]} \sum_{i=1}^n q^{v+\mu_i+n-i} \sum_{\sigma \in S_n} \epsilon(\sigma) \sigma x^{\mu+ve_i+\delta} \end{aligned}$$

This proves 4.3.

The next identity shows that R_v may be given an expression that is similar to the one obtained by Macdonald for his D_n^1 operator.

Proposition 4.2

We have

$$R_v = q^v \sum_{i=1}^n A_i(x; q) x_i^v T_{x_i}^q \quad 4.4$$

with

$$A_i(x; q) = \prod_{j \in [1, n]; j \neq i} \frac{q x_i - x_j}{x_i - x_j} . \quad 4.5$$

Proof

The definition in 4.1 may also be rewritten as

$$R_v P[X_n] = q^v \sum_{i=1}^n \left(\frac{1}{\Delta[X_n]} T_{x_i}^q \Delta[X_n] \right) x_i^v T_{x_i}^q P[X_n]$$

This shows 4.4 with

$$A_i(x, q) = \frac{1}{\Delta[X_n]} T_{x_i}^q \Delta[X_n].$$

However, we see that

$$\frac{1}{\Delta[X_n]} T_{x_i}^q \Delta[X_n] = \prod_{r, s \neq i} \frac{x_r - x_s}{x_r - x_s} \prod_{i < s} \frac{q x_i - x_s}{x_i - x_s} \prod_{r < i} \frac{x_r - q x_i}{x_r - x_i} = \prod_{j \in [1, n]; j \neq i} \frac{q x_i - x_j}{x_i - x_j},$$

as desired.

One of the difficulties in using Macdonald's original definition of the operators D_n^k stems from the fact that formulas expressing a symmetric polynomial $P[X_n]$ in terms of the variables themselves are quite impractical for significant values of n . For this reason, neither the definition in 4.1 nor its alternate form in 4.4 are much help in computing the action the operators D_n^k when it matters. However, for any of the operators R_v , we do have a plethystic formula which is computationally as well as theoretically very convenient. To state and prove this result we need some auxiliary identities

Proposition 4.3

For any formal series $F(x) = \sum_{k \geq 0} c_k x^k$ and for all integers $v \geq 0$ we have

$$F(1/z) \Omega[(q-1)zX_n] \Big|_{z^v} = \frac{\chi(v=0)}{q^n} F(0) + \frac{q-1}{q^n} \sum_{i=1}^n \left(\prod_{j=1; j \neq i}^n \frac{q x_i - x_j}{x_i - x_j} \right) (q x_i)^v F(q x_i) \quad 4.6$$

Proof

Starting from the partial fraction expansion

$$\Omega[(q-1)zX_n] = \prod_{i=1}^n \frac{1 - z x_i}{1 - z q x_i} = \frac{1}{q^n} + \frac{q-1}{q^n} \sum_{i=1}^n \left(\prod_{j=1; j \neq i}^n \frac{q x_i - x_j}{x_i - x_j} \right) \frac{1}{1 - z q x_i} ,$$

we get

$$\begin{aligned}
F(1/z) \Omega[(q-1)zX_n] \Big|_{z^v} &= \frac{F(1/z)}{q^n} \Big|_{z^v} + \frac{q-1}{q^n} \sum_{i=1}^n \left(\prod_{j=1; j \neq i}^n \frac{qx_i - x_j}{x_i - x_j} \right) \frac{1}{1 - zqx_i} F(1/z) \Big|_{z^v} \\
&= \frac{\chi(v=0)}{q^n} F(0) + \frac{q-1}{q^n} \sum_{i=1}^n \left(\prod_{j=1; j \neq i}^n \frac{qx_i - x_j}{x_i - x_j} \right) \left(\sum_{r \geq 0} (zqx_i)^r \right) \left(\sum_{s \geq 0} F_s / z^s \right) \Big|_{z^v} \\
&= \frac{\chi(v=0)}{q^n} F(0) + \frac{q-1}{q^n} \sum_{i=1}^n \left(\prod_{j=1; j \neq i}^n \frac{qx_i - x_j}{x_i - x_j} \right) \left(\sum_{r-s=v; r, s \geq 0} (qx_i)^r F_s \right) \\
&= \frac{\chi(v=0)}{q^n} F(0) + \frac{q-1}{q^n} \sum_{i=1}^n \left(\prod_{j=1; j \neq i}^n \frac{qx_i - x_j}{x_i - x_j} \right) \left(\sum_{s \geq 0} (qx_i)^{s+v} F_s \right)
\end{aligned}$$

This proves our Proposition.

As a corollary we obtain the following basic identity.

Theorem 4.1

For any symmetric polynomial $P[X_n]$ we have

$$R_v P[X_n] = \frac{\chi(v=0)}{1-q} P[X_n] + \frac{q^{n+v}}{q-1} P[X_n - (1-q)/z] \Omega[(1-1/q)zX_n] \Big|_{z^v}. \quad 4.7$$

Proof

Notice that in view of 4.4 we can rewrite 4.1 in the form

$$\begin{aligned}
R_v P[X_n] &= q^v \sum_{i=1}^n A_i(x; q) x_i^v P[X_n - (1-q)x_i] \\
&= q^v \sum_{i=1}^n \left(\prod_{j=1, j \neq i}^n \frac{qx_i - x_j}{x_i - x_j} \right) x_i^v P[X_n - (1-q)x_i].
\end{aligned} \quad 4.8$$

On the other hand Proposition 4.3 for $F(z) \rightarrow F(z/q)$ yields

$$F(1/qz) \Omega[(q-1)zX_n] \Big|_{z^v} = \frac{\chi(v=0)}{q^n} F(0) + \frac{q-1}{q^n} \sum_{i=1}^n \left(\prod_{j=1; j \neq i}^n \frac{qx_i - x_j}{x_i - x_j} \right) (qx_i)^v F(x_i)$$

Or better

$$q^v \sum_{i=1}^n \left(\prod_{j=1, j \neq i}^n \frac{qx_i - x_j}{x_i - x_j} \right) x_i^v F(x_i) = \frac{\chi(v=0)}{1-q} F(0) + \frac{q^n}{q-1} F(1/qz) \Omega[(q-1)zX_n] \Big|_{z^v}.$$

Using this with $F(z) = P[X_n - (1-q)z]$ and using 4.8 gives

$$\begin{aligned}
R_v P[X_n] &= q^v \sum_{i=1}^n \left(\prod_{j=1, j \neq i}^n \frac{qx_i - x_j}{x_i - x_j} \right) x_i^v P[X_n - (1-q)x_i] \\
&= \frac{\chi(v=0)}{1-q} P[X_n] + \frac{q^n}{q-1} P[X_n - (1-q)/qz] \Omega[(1-1/q)qzX_n] \Big|_{z^v}.
\end{aligned} \quad 4.9$$

Notice next that for any two formal power series $A(z), B(z)$ we have the identity

$$\begin{aligned}
A[1/qz] B[qz] \Big|_{z^v} &= \sum_{r,s} A_r B_s \left(\frac{1}{qz} \right)^r (qz)^s \Big|_{z^v} = \sum_{r,s} (zq)^{s-r} A_r B_s \Big|_{z^v} \\
&= \sum_{s-r=v} q^v A_r B_s = q^v \sum_r A_r B_{r+v} = q^v A[1/z] B[z] \Big|_{z^v}
\end{aligned}$$

and thus 4.9 becomes

$$R_v P[X_n] = \frac{\chi(v=0)}{1-q} P[X_n] + \frac{q^{n+v}}{q-1} P[X_n - (1-q)/z] \Omega[(1-1/q)zX_n] \Big|_{z^v}.$$

This proves 4.7.

In particular, setting $P[X_n] = s_\mu[X_n]$ and using 4.3 we obtain

$$\sum_{i=1}^n q^{v+\mu_i+n-i} s_{\mu+ve_i}[X_n] = \frac{\chi(v=0)}{1-q} s_\mu[X_n] + \frac{q^{n+v}}{q-1} s_\mu[X_n - (1-q)/z] \Omega[(1-1/q)zX_n] \Big|_{z^v}.$$

or better yet

$$q^{n+v} s_\mu[X_n - (1-q)/z] \Omega[(1-1/q)zX_n] \Big|_{z^v} = \chi(v=0) s_\mu[X_n] + (q-1) \sum_{i=1}^n q^{v+\mu_i+n-i} s_{\mu+ve_i}[X_n] \quad 4.10$$

At this point it is more convenient to separate the cases $v > 0$ and $v = 0$. We will begin with the following immediate corollary of Theorem 4.1.

Proposition 4.4

For any $u, v > 0$ and any partition μ we have

$$q^{uv} s_\mu[X - (1-q^u)/z] \Omega[(1-q^{-u})zX] \Big|_{z^v} = (q^u - 1) \sum_{i=1}^{|\mu|+v} q^{u(p(\mu)_i+v-i)} s_{p(\mu)+ve_i}[X] \quad 4.11$$

where $p(\mu)$ is the weak composition of length $|\mu| + v$ obtained by adjoining zeros to the parts of μ and e_i is the i^{th} coordinate vector of length $|\mu| + v$.

Proof

For $v > 0$ 4.10 can be rewritten in the form

$$s_\mu[X_n - (1-q)/z] \Omega[(1-1/q)zX_n] \Big|_{z^v} = (q-1) \sum_{i=1}^n q^{\mu_i+v-i} s_{\mu+ve_i}[X_n]$$

and the replacement $q \rightarrow q^u$ gives

$$q^{uv} s_\mu[X_n - (1-q^u)/z] \Omega[(1-q^{-u})zX_n] \Big|_{z^v} = (q^u - 1) \sum_{i=1}^n q^{u(\mu_i+v-i)} s_{p(\mu)+ve_i}[X_n]$$

This given, 4.11 follows since the Schur functions involved in this expression stabilize after $n \geq |\mu| + v$.

Keeping this in mind let us recall that our goal here is to work out the action of the operator $D_{u,v}$ on the basis $\{s_\mu[\frac{X}{1-q}]\}_\mu$. The following identity provides the link that ties this goal with the identities in 4.7 and 4.11

Proposition 4.5

Suppose that for some $v \geq 0$ we have

$$s_\mu[X - (1-q^u)/z] \Omega[(1-q^{-u})zX] \Big|_{z^v} = G[X; q] \quad 4.12$$

then

$$D_{u,v} s_\mu[\frac{X}{1-q}] = q^v G[\frac{X}{1-q}; q] \quad 4.13$$

Proof

Notice first the following sequence of equalities

$$\begin{aligned}
q^v G[X; q] &= q^v s_\mu [X - (1 - q^u)/z] \Omega[z(1 - q^{-u})X] \Big|_{z^v} \\
&= s_\mu [X - (1 - q^u)/qz] \Omega[qz(1 - q^{-u})X] \Big|_{z^v} \\
&= s_\mu [X + \frac{(q-1)}{1-q} (1 - q^u)/qz] \Omega[-qz \frac{1-q^{-u}}{q-1} X(1 - q)] \Big|_{z^v} \\
&= s_\mu [X + \frac{(1-1/q)}{1-q} (1 - q^u)/z] \Omega[-z \frac{1-q^{-u}}{1-q^{-1}} X(1 - q)] \Big|_{z^v}
\end{aligned}$$

Next the replacement $X \rightarrow \frac{X}{1-q}$ gives

$$\begin{aligned}
q^v G[\frac{X}{1-q}; q] &= s_\mu [\frac{X}{1-q} + \frac{(1-1/q)}{1-q} (1 - q^u)/z] \Omega[-z \frac{1-q^{-u}}{1-q^{-1}} X] \Big|_{z^v} \\
&= s_\mu [\frac{X + (1-1/q)(1-q^u)/z}{1-q}] \Omega[-z \frac{1-q^{-u}}{1-q^{-1}} X] \Big|_{z^v} \\
&= s_\mu [\frac{X + (1-q)(1-1/q)[u]_q/z}{1-q}] \Omega[-z \frac{1-q^{-u}}{1-q^{-1}} X] \Big|_{z^v}
\end{aligned} \tag{4.14}$$

Now recalling that by definition we have (for $t = 1/q$)

$$D_{u,v} F[X] = F[X + M[u]_q/z] \Omega[-z[u]_t X] \Big|_{z^v},$$

we see that 4.14 proves 4.13.

We are thus able to obtain our

Proof of Theorem I.6

By combining Propositions 4.4 and 4.5 with

$$G[X; q] = q^{-uv} (q^u - 1) \sum_{i=1}^{|\mu|+v} q^{u(p(\mu)_i + v - i)} s_{p(\mu) + v e_i} [X]$$

we obtain

$$D_{u,v} s_\mu [\frac{X}{1-q}] = (q^u - 1) \sum_{i=1}^{|\mu|+v} q^{u p(\mu)_i + v - u i} s_{p(\mu) + v e_i} [\frac{X}{1-q}]$$

as desired.

Our next task is to take care of the case $v = 0$ of 4.7. This may be rewritten as

$$q^n P[X_n - (1 - q)/z] \Omega[1 - 1/q] z X_n \Big|_{z^0} = P[X_n] - (1 - q) R_0 P[X_n]$$

Choosing $P[X_n] = s_\mu[X_n]$ and using 4.3 for $v = 0$ we get

$$q^n s_\mu [X_n - (1 - q)/z] \Omega[1 - 1/q] z X_n \Big|_{z^0} = \left(1 - (1 - q) \sum_{i=1}^n q^{\mu_i + n - i}\right) s_\mu[X_n] \tag{4.15}$$

Our next step is to transform 4.15 into a relation which contains no explicit dependence on n . To this end, recalling that we write a partition of n in the form $\mu = (\mu_1 \geq \mu_2 \geq \dots \geq \mu_n \geq 0)$ we set

$$B_\mu(q, t) = \sum_{\mu_i > 0} t^{i-1} (1 + q + \dots + q^{\mu_i - 1}) = \sum_{i=1}^n t^{i-1} \frac{1 - q^{\mu_i}}{1 - q} \tag{4.16}$$

This polynomial, usually called the *bisexponent generator* of μ , plays an essential role in the Theory of Macdonald polynomials, in our case this results in the following basic identity:

Proposition 4.6

$$s_\mu[X_n - \frac{1-q}{z}] \Omega[(1-q^{-1})zX_n] \Big|_{z_0} = \left(1 - (1-q^{-1})(1-q)B_\lambda(q, q^{-1})\right) s_\mu[X_n] \quad 4.17$$

Proof

Changing t into $1/q$ in 4.16 and multiplying both sides by q^n we can rewrite it in the form

$$q^n \left(1 - (1-q^{-1})(1-q)B_\mu(q, t)\right) = 1 - (1-q) \sum_{i=1}^n q^{\mu_i+n-i} \quad 4.18$$

Using this in 4.15 gives

$$q^n s_\mu[X_n - \frac{1-q}{z}] \Omega[(1-q^{-1})zX_n] \Big|_{z_0} = q^n \left(1 - (1-q^{-1})(1-q)B_\mu(q, t)\right) s_\mu[X_n]$$

canceling the factor q^n proves 4.17 as desired.

Remark 4.1

By setting $v = 0$ in 4.3 and 4.4 we derive that

$$\sum_{i=1}^n A_i(x; q) T_{x_i}^q s_\mu[X_n] = \left(\sum_{i=1}^n q^{\mu_i+n-i}\right) s_\mu[X_n]$$

This identity was used by Macdonald in [25] to prove that his polynomial $P_\mu(X_n, q, t)$ reduces to $s_\mu[X_n]$ at $t = q$. Now the original definition of the modified Macdonald polynomial $\tilde{H}_\mu[X; q, t]$ was obtained by setting

$$\tilde{H}_\mu[X; q, t] = c_\nu(q, t) P_\mu[X/(1-1/t); q, 1/t] \quad 4.19$$

where $P_\mu[X; q, t]$ is none other than $P_\mu[X_n; q, t]$ (for $\mu \vdash n$) with X_n replaced by the infinite alphabet $X = x_1 + x_2 + \dots$ and $c_\nu(q, t)$ is a polynomial in q, t whose nature is immaterial here. Thus it follows from 4.19 that

$$\tilde{H}_\mu[X; q, 1/q] = c_\nu(q, q^{-1}) P_\mu[X/(1-q); q, q] = c_\nu(q, q^{-1}) s_\mu\left[\frac{X}{1-q}\right]$$

Thus for all practical purposes, in the present context, which arises from our setting $t = 1/q$ in all our $Q_{u,v}$ and their identities, the basis $\{\tilde{H}_\mu[X; q, t]\}_\mu$ need only be replaced by the basis $\{s_\mu[\frac{X}{1-q}]\}_\mu$. In this vein we can easily obtain an alternate way of interpreting the identity in 4.17.

Theorem I.5

For all integers $u \geq 1$ we have

$$D_{u,0} s_\mu\left[\frac{X}{1-q}\right] = \left(1 - (1-q^{-u})(1-q^u)B_\mu(q^u, q^{-u})\right) s_\mu\left[\frac{X}{1-q}\right] \quad 4.20$$

Proof

Making the replacements $X_n \rightarrow X$ and $q \rightarrow q^u$ in 4.16 gives

$$s_\mu\left[X - \frac{1-q^u}{z}\right] \Omega[(1-q^{-u})zX] \Big|_{z_0} = \left(1 - (1-q^{-u})(1-q^u)B_\mu(q^u, q^{-u})\right) s_\mu[X]$$

Next we do $X \rightarrow \frac{X}{1-q}$ and get

$$s_\mu\left[\frac{X}{1-q} - \frac{1-q^u}{z}\right] \Omega\left[-\frac{1-q^{-u}}{1-1/q}(z/q)X\right] \Big|_{z_0} = \left(1 - (1-q^{-u})(1-q^u)B_\mu(q^u, q^{-u})\right) s_\mu\left[\frac{X}{1-q}\right]$$

and this (with $t = 1/q$) may be rewritten as

$$s_\mu \left[\frac{X + (1-1/q)(1-q)[m]_q/(z/q)}{1-q} \right] \Omega[-[m]_t(z/q)X] \Big|_{z^0} = \left(1 - (1 - q^{-u})(1 - q^u)B_\mu(q^u, q^{-u}) \right) s_\mu \left[\frac{X}{1-q} \right]$$

or better

$$s_\mu \left[\frac{X + M[m]_q/z}{1-q} \right] \Omega[-[m]_t z X] \Big|_{z^0} = \left(1 - (1 - q^{-u})(1 - q^u)B_\mu(q^u, q^{-u}) \right) s_\mu \left[\frac{X}{1-q} \right] \quad 4.21$$

Recalling that by definition we have

$$D_{u,v}F[X] = F[X + M[u]_q/z] \Omega[-[u]_t z X] \Big|_{z^v}.$$

We see that 4.20 is simply another way of writing 4.21. This completes our proof.

Remark 4.2

If we follow the sequence of steps that yielded the identity in 4.20 we will notice that this identity is but a direct consequence of the identity in 4.17 with the replacement $X_n \rightarrow X$, that is

$$s_\mu \left[X - \frac{1-q}{z} \right] \Omega[(1 - q^{-1})zX] \Big|_{z^0} = \left(1 - (1 - q^{-1})(1 - q)B_\mu(q, q^{-1}) \right) s_\mu[X]. \quad 4.22$$

Thus in principle, a short cut in the proof of Theorem I.5, may appear to be the verification of 4.22.

To better appreciate the power of the path we followed in the proof of 4.20, it will be instructive to see what kind of combinatorial identities we are led to in trying to carry this out. Now working first on the left hand side of 4.22 gives

$$\begin{aligned} LHS &= s_\mu[X] + \sum_{k \geq 1} \sum_{\nu=(k-a, 1^a)} s_{\mu/\nu'}[X] (-1/z)^k s_\nu[1-q] \Omega[(1 - q^{-1})zX] \Big|_{z^0} \\ &= s_\mu[X] + (1-q) \sum_{k \geq 1} (-1)^k \sum_{a=0}^{k-1} s_{\mu/(a+1, 1^{k-a-1})}[X] (-q)^a h_k[1 - q^{-1}]X \end{aligned} \quad 4.23$$

and since

$$h_k[1 - q^{-1}]X = (1 - q^{-1}) \sum_{b=0}^{k-1} (-q)^b s_{k-b, 1^b}[X]$$

4.23 becomes

$$LHS = s_\mu[X] + (1-q)(1 - q^{-1}) \sum_{k \geq 1} (-1)^k \sum_{a,b=0}^{k-1} (-q)^{a+b} s_{\mu/(a+1, 1^{k-a-1})}[X] s_{k-b, 1^b}[X] \quad 4.24$$

Taking the scalar product of both sides of 4.22 by $s_\lambda[X]$ and using 4.24, routine manipulations reduce 4.22 to the equivalent identity

$$\sum_{k \geq 1} (-1)^k \sum_{a,b=0}^{k-1} (-q)^{a+b} \langle s_{\mu/(a+1, 1^{k-a-1})}, s_{\lambda/(k-b, 1^b)} \rangle = -\chi(\lambda = \mu) B_\mu(q, q^{-1}) \quad 4.25$$

A standard result on scalar products of skew Schur functions (see [13]), asserts that the scalar product summand is none other than the number of permutations that fit the shape $\mu/(a+1, 1^{k-a-1})$ whose inverse fits the shape $\lambda/(k-b, 1^b)$. This given, the only conclusion we can draw from this calculation is that 4.25 is one truly remarkable combinatorial consequence of Theorem I.5.

5. The original proof of Theorem I.1 by the partial fraction method

In this section we explain how we discovered Theorem I.1. Indeed Theorem I.1 involves different series expansions of a single rational function, which is best understood by using the partial fraction method of the fourth named author. To this end we need to work in the field $K = \mathbb{Q}((z_N))((z_{N-1})) \cdots ((z_1))$ of iterated Laurent series to obtain series expansion of rational functions. The readers are referred to [31] for the original development of the field of iterated Laurent series. Here we only recall that K defines a total group order on its monomials given by

$$z_1^{a_1} \cdots z_N^{a_N} \begin{cases} <_K 1, & \text{if } a_1 = \cdots = a_{i-1} = 0 \text{ \& } a_i > 0; \\ = 1, & \text{if } a_1 = \cdots = a_N = 0; \\ >_K 1, & \text{if } a_1 = \cdots = a_{i-1} = 0 \text{ \& } a_i < 0. \end{cases}$$

We shall simply write this order by $z_1 < z_2 < \cdots < z_N < 1$. The series expansion of $(1 - w)^{-1}$ for a monomial $w \neq 1$ (called small or large) is thus given by

$$\frac{1}{1 - w} = \begin{cases} \sum_{n \geq 0} w^n, & \text{if } w <_K 1; \\ \frac{1}{-w(1-1/w)} = -\sum_{n \geq 0} w^{-n-1}, & \text{if } w >_K 1. \end{cases}$$

To start with, let us recall that for any Laurent polynomials $L(z_1, z_2) \in \mathbb{Q}[z_1^{\pm 1}, z_2^{\pm 1}]$, we always have that

$$L(z_1, z_2) \Big|_{z_1^0 z_2^0} = L(z_2, z_1) \Big|_{z_1^0 z_2^0} \quad \text{holds in } \mathbb{Q}[z_1^{\pm 1}, z_2^{\pm 1}].$$

In particular, antisymmetric Laurent polynomials have constant term 0. Such properties no longer hold for rational functions in $K_1 = \mathbb{Q}((z_1))((z_2))$. For example, $f(z_1, z_2) = \frac{z_1 + z_2}{z_1 - z_2}$ is clearly antisymmetric, but in K_1 we have

$$\frac{z_1 + z_2}{z_1 - z_2} \Big|_{z_1^0 z_2^0} = \frac{1 + z_2/z_1}{1 - z_2/z_1} \Big|_{z_1^0 z_2^0} = (1 + z_2/z_1) \sum_{n \geq 0} (z_2/z_1)^n \Big|_{z_1^0 z_2^0} = 1.$$

Indeed, the exchanging of the two variables z_1, z_2 transforms the constant term in K_1 to a constant term in $K_2 = \mathbb{Q}((z_2))((z_1))$. We shall have

$$f(z_1, z_2) \Big|_{z_1^0 z_2^0}^{K_1} = f(z_2, z_1) \Big|_{z_1^0 z_2^0}^{K_2}$$

where the left hand side is a constant term in K_1 , but the right hand side is a constant term in K_2 . This type of exchanging of variables would be sufficient for us to prove Theorem I.1. See [30] for general formulation on the change of variables in a field of Malcev-Neumann series.

Proof of Theorem I.1

With $t = 1/q$, we have

$$\begin{aligned} D_{c,d} D_{a,b} F[X] &= D_{c,d} F[X + [a]_q \frac{M}{z_1}] \Omega[-z_1 X[a]_t] z_1^{-b} \Big|_{z_1^0} \\ &= F[X + [c]_q \frac{M}{z_2} + [a]_q \frac{M}{z_1}] \Omega[-z_1[a]_t(X + [c]_q \frac{M}{z_2})] \Omega[-z_2 X[c]_t] z_1^{-b} z_2^{-d} \Big|_{z_1^0 z_2^0} \\ &= F[X + [c]_q \frac{M}{z_2} + [a]_q \frac{M}{z_1}] \Omega[-X(z_1[a]_t + z_2[c]_t)] \Omega[-M[a]_t[c]_q \frac{z_1}{z_2}] z_1^{-b} z_2^{-d} \Big|_{z_1^0 z_2^0} \\ &= F[X + [c]_q \frac{M}{z_2} + [a]_q \frac{M}{z_1}] \Omega[-X(z_1[a]_t + z_2[c]_t)] \frac{(1 - \frac{z_1}{z_2})(1 - q^c t^a \frac{z_1}{z_2})}{(1 - q^c \frac{z_1}{z_2})(1 - t^a \frac{z_1}{z_2})} z_1^{-b} z_2^{-d} \Big|_{z_1^0 z_2^0} \end{aligned}$$

where in the last step we have used the fact that $M[a]_t[c]_q = (1 - t^a)(1 - q^c) = 1 - t^a - q^c + q^c t^a$. This constant term has to be understood as in a field of iterated Laurent series where $q^c z_1/z_2$ and $t^a z_1/z_2$ are

small. In the general q, t case, we can set $q < t < z_1 < z_2 < 1$. But here we can not set q to be small, since that will force $t = 1/q$ to be large. We choose to work in the field of iterated Laurent series K_1 defined by the order $z_1 < z_2 < q < 1$ (one can take q as a constant). More precisely, we can set, for example, $K_1 = \mathbb{Q}(q)((z_2))((z_1))[[x_1, x_2, \dots]]$.

Let us write

$$D_{c,d}D_{a,b}F[X] = F \left[X + [c]_q \frac{M}{z_2} + [a]_q \frac{M}{z_1} \right] \Omega [-X(z_1[a]_t + z_2[c]_t)] G(z_1, z_2; c, d, a, b) \Big|_{z_1^0 z_2^0}^{K_1}, \quad 5.1$$

where

$$G(z_1, z_2; c, d, a, b) = \frac{(1 - \frac{z_1}{z_2})(1 - q^c t^a \frac{z_1}{z_2})}{(1 - q^c \frac{z_1}{z_2})(1 - t^a \frac{z_1}{z_2})} z_1^{-b} z_2^{-d}.$$

Now switching (c, d) and (a, b) gives

$$D_{a,b}D_{c,d}F[X] = F \left[X + [a]_q \frac{M}{z_2} + [c]_q \frac{M}{z_1} \right] \Omega [-X(z_1[c]_t + z_2[a]_t)] G(z_1, z_2; a, b, c, d) \Big|_{z_1^0 z_2^0}^{K_1}.$$

Observe that when $t = 1/q$, we have

$$G(z_2, z_1; a, b, c, d) = \frac{(1 - \frac{z_2}{z_1})(1 - t^c q^a \frac{z_2}{z_1})}{(1 - t^c \frac{z_2}{z_1})(1 - q^a \frac{z_2}{z_1})} = \frac{(1 - \frac{z_1}{z_2})(1 - q^c t^a \frac{z_1}{z_2})}{(1 - q^c \frac{z_1}{z_2})(1 - t^a \frac{z_1}{z_2})} = G(z_1, z_2; c, d, a, b).$$

By exchanging z_1 and z_2 , we obtain that $D_{a,b}D_{c,d}F[X]$ is the same constant term as in 5.1, but working in the field of iterated Laurent series K_2 defined by the order $z_2 < z_1 < q < 1$. So we are indeed computing the difference of the constant terms of a single “rational function” in two different working fields.

By partial fraction decomposition in z_1 , applied to the coefficients in the x ’s and then sum, we have

$$\begin{aligned} F \left[X + [c]_q \frac{M}{z_2} + [a]_q \frac{M}{z_1} \right] \Omega [-X(z_1[a]_t + z_2[c]_t)] G(z_1, z_2; a, b, c, d) = \\ = p^{\geq 0}(z_1) + p^{< 0}(z_1) + \frac{A_1}{1 - q^c \frac{z_1}{z_2}} + \frac{A_2}{1 - t^a \frac{z_1}{z_2}}, \end{aligned}$$

where when we restrict to each coefficient of the x ’s, $p^{\geq 0}(z_1)$ is a polynomial in z_1 , $p^{< 0}(z_1)$ is a Laurent polynomial that only contains negative powers in z_1 , and A_1 and A_2 are free of z_1 given by

$$\begin{aligned} A_1 &= F \left[X + [c]_q \frac{M}{z_2} + q^c [a]_q \frac{M}{z_2} \right] \Omega [-X(z_2 t^c [a]_t + z_2 [c]_t)] \frac{(1 - t^c)(1 - t^a)}{(1 - t^a t^c)} z_2^{-b} t^{-bc} z_2^{-d}, \\ &= -q^{bc} \frac{(1 - q^c)(1 - q^a)}{(1 - q^{a+c})} F \left[X + [a + c]_q \frac{M}{z_2} \right] \Omega [-X z_2 [a + c]_t] z_2^{-b-d}, \\ A_2 &= F \left[X + [c]_q \frac{M}{z_2} + t^a [a]_q \frac{M}{z_2} \right] \Omega [-X(z_2 q^a [a]_t + z_2 [c]_t)] \frac{(1 - q^a)(1 - q^c)}{(1 - q^a q^c)} z_2^{-b} q^{-ba} z_2^{-d} \\ &= \frac{(1 - q^a)(1 - q^c)}{(1 - q^a q^c)} F \left[X + [a + c]_q \frac{M}{z_2 q^a} \right] \Omega [-X z_2 q^a [a + c]_t] z_2^{-d-b} q^{-ba}. \end{aligned} \quad 5.2$$

Actually we need the following formula (obtained by the change of variables $z_2 \rightarrow z_2 t^a$).

$$A_2 \Big|_{z_2^0} = q^{da} \frac{(1 - q^a)(1 - q^c)}{(1 - q^{a+c})} F \left[X + [a + c]_q \frac{M}{z_2} \right] \Omega [-X z_2 [a + c]_t] z_2^{-d-b} \Big|_{z_2^0}. \quad 5.3$$

Now we take the constant term in z_1 first, working in K_1 and K_2 separately. Since the two monomials $q^c z_1/z_2$ and $t^a z_1/z_2$ are both small in K_1 but large in K_2 , we have

$$\begin{aligned} D_{c,d}D_{a,b}F[X] &= p^{\geq 0}(0) \Big|_{z_2^0} + A_1 \Big|_{z_2^0} + A_2 \Big|_{z_2^0}. \\ D_{a,b}D_{c,d}F[X] &= p^{\geq 0}(0) \Big|_{z_2^0}. \end{aligned}$$

It follows that

$$\frac{1}{M}[D_{c,d}, D_{a,b}]F[X] = \frac{1}{M}\left(A_1\Big|_{z_2^0} + A_2\Big|_{z_2^0}\right).$$

Applying formulas 5.2 and 5.3 gives the desired result.

The idea of the proof of Theorem I.1 can be generalized as follows, which is easy to prove but turns out to be very useful.

Proposition 5.1

Let K_1 and K_2 be two different field of iterated Laurent series. Suppose that we have the following partial fraction expansion:

$$F(z) = p_0(z) + \frac{p_{-1}(z)}{z^m} + \frac{p_1(z)}{(1 - u_1 z)^{k_1}} + \cdots + \frac{p_N(z)}{(1 - u_N z)^{k_N}}.$$

Then we have

$$F(z)\Big|_{z^0}^{K_1} - F(z)\Big|_{z^0}^{K_2} = \sum_{u_i z <_{K_1} 1 \text{ \& } u_i z >_{K_2} 1} p_i(0) - \sum_{u_j z >_{K_1} 1 \text{ \& } u_j z <_{K_2} 1} p_j(0).$$

In words, the difference of the two constant terms only came from those denominators that are contributing (i.e., with $u_i z < 1$) in one field but dually contributing (i.e., with $u_i z > 1$) in the other field.

Proof

With the given partial fraction decomposition, taking constant term in z under K_1 gives

$$F(z)\Big|_{z^0}^{K_1} = p_0(0) + \sum_{u_i z^{k_i} <_{K_1} 1} p_i(0).$$

A similar result holds for K_2 . Subtracting gives the desired formula.

Remark: The proposition applies if $F(z)$ includes something like $\Omega[X + M/z]$ or $\Omega[-zX]$ as factors, in which case m or $p_0(z)$ does not exist. But $F(z)$ can be first expanded as a power series in the x 's, and then apply the proposition to the coefficients in the x 's.

Remark: If we take $K_1 = \mathbf{C}((z))$ and $K_2 = \mathbf{C}((z^{-1}))$, then the proposition gives

$$F(z)\Big|_{z^0}^{K_1} - F(z)\Big|_{z^0}^{K_2} = \sum_i p_i(0).$$

This can be shown to be equivalent to the well-known fact that for any given rational function, its residues at all points (including ∞) sum to 0.

Bibliography

- [1] F. Bergeron, A. M. Garsia, M. Haiman and G. Tesler, *Identities and positivity conjectures for some remarkable operators in the theory of symmetric functions*, Methods Appl. Anal. 6 (1999), 363-420.
- [2] F. Bergeron and A. M. Garsia, *Science Fiction and Macdonald's Polynomials*. CRM Proceedings & Lecture Notes, American Mathematical Society, 22: 1–52, 1999.
- [3] F. Bergeron, A. Garsia, E. Leven, G. Xin, *Compositional (km, kn) -Shuffle Conjectures*, math arXiv:1404.4616.
- [4] F. Bergeron, A. M. Garsia, E. Leven and G. Xin, *Some remarkable new Plethystic operators in the Theory of Macdonald Polynomials*, math arXiv: 1405.0316.
- [5] I. Burban and O. Schiffmann, *On the Hall algebra of an elliptic curve, I*, Preprint (2005), arXiv:math.AG/0505148.
- [6] C. Dunkl, *Intertwining operators and polynomials associated with the symmetric group*, Monatsh. Math. 126 (1998), no. 3, 181-209.
- [7] E. Egge, N. Loehr and G. Warrington, *From quasisymmetric expansions to Schur expansions via a modified inverse Kostka matrix*, European Journal of Combinatorics, V. 1. Dec 2010 pp. 2014-2027
- [8] A. M. Garsia and J. Haglund, *Proof of the q, t -Catalan Positivity Conjecture*, Discrete Mathematics, V 256, Issue 3, 28 October 2002, pp. 677-717
- [9] A.M. Garsia, J. Haglund, G. Xin, M. Zabrocki *Some new applications of the Stanley-Macdonald Pieri Rules*, arXiv:1407.7916, Comments: for Stanley@70. Massachusetts institute of technology, June 23-27, 2014
- [10] A. Garsia and M. Haiman, *A remarkable q, t -Catalan sequence and q -Lagrange inversion*, J. Algebraic Combin. 5 (1996), no. 3, 191-244.
- [11] A. M. Garsia and M. Haiman, *Some Natural Bigraded S_n -Modules and q, t -Kostka Coefficients. (1996) (electronic) The Foata Festschrift*, [http://www.combinatorics.org/Volume 3/volume 3 2.html#R24](http://www.combinatorics.org/Volume%203/volume%203%20.html#R24)
- [12] A. M. Garsia, M. Haiman, and G. Tesler. *Explicit plethystic formulas for Macdonald (q, t) -Kostka coefficients*. Séminaire Lotharingien de Combinatoire [electronic only], 42:B42m, 1999.
- [13] A. Garsia and J. Remmel, *Shuffles of permutations and the Kronecker product*, Graphs and Combinatorics, V 1, issue 1, (1985) pp. 217-263.
- [14] A. M. Garsia, G. Xin and M. Zabrocki, *Hall-Littlewood Operators in the Theory of Parking Functions and Diagonal Harmonics*, Int Math Res Notices (2011) doi: 10.1093/imrn/rnr060
- [15] I. Gessel. *Multipartite P -partitions and inner products of skew Schur functions*. Contemp. Math, 34: 289–301, 1984.
- [16] E. Gorsky, *Arc spaces and DAHSA representations*, math arXiv:1110.1674 (2011).
- [17] E. Gorsky and M. Mazin, *Compactified Jacobians and q, t -Catalan Numbers I*, Journal of Combinatorial Theory, Series A, 120 (2013) 49-63.
- [18] , E. Gorsky and M. Mazin, *Compactified Jacobians and q, t -Catalan Numbers II*, Journal of Alg, Comb. I (2014) **39**: 153-186.
- [19] E. Gorsky and A. Negut. *Refined knot invariants and Hilbert schemes*. math arXiv:1304.3328, 2013.
- [20] J. Haglund, M. Haiman, N. Loehr, J. B. Remmel, and A. Ulyanov. *A combinatorial formula for the character of the diagonal coinvariants*. Duke J. Math., 126:195-232, 2005.
- [21] J. Haglund, J. Morse, and M. Zabrocki. *A compositional refinement of the shuffle conjecture specifying touch points of the Dyck path*. Canadian J. Math, 64: 822-844, 2012.

- [22] M. Haiman, *Conjectures on the Quotient Ring by Diagonal Invariants*, Journal of Alg. Comb. **3** (1994), 17–76.
- [23] A. Hicks and E. Leven. *A simpler formula for the number of diagonal inversions of an (m,n) -Parking Function and a returning Fermionic formula*, math arXiv:1404.4889 (2014): (to appear in Discrete Math.)
- [24] T. Hikita. *Affine springer fibers of type A and combinatorics of diagonal coinvariants*. math arXiv:1203.5878, 2012.
- [25] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Mathematical Monographs, Oxford Science Publications, Oxford University Press, New York, 1995.
- [26] O. Schiffmann and E. Vasserot. *The elliptical Hall algebra, Cherednik Hecke algebras and Macdonald polynomials*. Compos. Math., 147.1:188–234, 2011.
- [27] O. Schiffmann and E. Vasserot. *The elliptical Hall algebra and the equivariant K-theory of the Hilbert scheme of A^2* . Duke J. Math., 162.2: 279–366, 2013.
- [28] O. Schiffmann, *On the Hall algebra of an elliptic curve, II*, Preprint (2005), arXiv:math/0508553.
- [29] S. Stevan. Chern-Simons invariants of torus links. Ann. Henri Poincare 11 (2010), no. 7, 1201–1224.
- [30] G. Xin, *A residue theorem for Malcev-Neumann series*, Adv. Appl. Math. (2005) 35, 271–293.
- [31] G. Xin, *A fast algorithm for MacMahon’s partition analysis*, Electron. J. Combin., (2004), 11, R53.